

Normalization of Polynomials in Algebraic Invariants of Three-Dimensional Orthogonal Geometry

Hongbo Li

KLMM, AMSS, Chinese Academy of Sciences, Beijing 100190, China

hli@mmrc.iss.ac.cn

ABSTRACT

In classical invariant theory, the Gröbner base of the ideal of syzygies and the normal forms of polynomials of invariants are two core contents. To improve the performance of invariant theory in symbolic computing of classical geometry, advanced invariants are introduced via Clifford product [5]. This paper addresses and solves the two key problems in advanced invariant theory: the Gröbner base of the ideal of syzygies among advanced invariants, and the normal forms of polynomials of advanced invariants. These results beautifully extend the straightening of Young tableaux to advanced invariants.

Keywords

Invariant theory; Clifford algebra; bracket algebra; non-commutative Gröbner base; straightening of Young tableaux.

1. INTRODUCTION

In traditional analytic approach to classical geometry, coordinates are introduced to represent points in the geometric space, and equations of the coordinates are used to define constraints among the points, forming a representation of higher dimensional objects such as curves, surfaces, etc. Basic manipulations of coordinates include addition and multiplication, resulting in polynomials in the coordinates. Since the coordinates of generic points are independent, and the multiplication of coordinate variables are commutative, normalization of the polynomials in the coordinates is very easy. The normal forms of the polynomials are required in many manipulations, *e.g.*, division among polynomials.

Another analytic approach to classical geometry, dating back to Euclid, is to use geometric invariants such as lengths, angles, areas, *etc.* A typically algebraic system of geometric invariants is a polynomial ring generated by basic invariants. In such a system, a vector variable in a linear space is used to represent a point or direction in classical geometry, the inner product of a vector with itself represents the squared

length of the vector, the inner product of two unit vectors represents the cosine of the angle between them, *etc.* Such operators among vectors generate a set of *basic invariants*, and the polynomials in these basic invariants are *advanced invariants*.

Although the multiplication of invariants are commutative, the basic invariants generated by generic vector variables of the linear space are not independent, and there are polynomial relations among them, called *syzygy* relations. The dependency is largely caused by the dimension constraint of the linear space upon vectors. While the dimension constraint can be easily reflected by the number of coordinates introduced to represent a point and the independency among the coordinates, for basic invariants generated by the points, fully representing the dimension constraint is by no means trivial. Classical invariant theory studies the generators of invariants, the syzygy relations among the basic invariants, and the normal forms of advanced invariants as polynomials in the basic ones [11], [12].

In symbolic geometric computing, both the coordinate approach and the basic invariant approach encounter the difficulty of very big polynomial size, in particular in the middle of symbolic manipulations. In [5], a recipe to alleviate the difficulty is proposed, called long geometric product, BREEFS, and Clifford factorization, among which the long geometric product (or Clifford product) is the foundation. The idea is to convert polynomials of basic invariants into advanced invariants, converse to the approach of classical invariant theory, by means of an associative and multilinear product among the vector variables representing points. The associativity of the product and the symmetries within a long bracket provide powerful manipulations that cannot be done with basic invariants, nor with coordinates. This is a top-down approach to advanced invariants [6], while the classical invariant theory is a bottom-up approach.

Dealing with the syzygy relations among advanced invariants and finding the normal forms of polynomials in advanced invariants are two fundamental tasks in such “advanced invariant theory”. The 2D case is easy, while higher dimensional cases are difficult. Little advance has been achieved in six years since the publication of [5] in 1997.

In this paper, the two fundamental problems are solved for the advanced invariant theory of 3D orthogonal geometry: the Gröbner base of the syzygy ideal of “long brackets”, and

the normal forms of Clifford bracket polynomials. It turns out that the normal forms of such bracket polynomials are surprisingly “beautiful”. The description is the following.

In classical invariant theory for $(n-1)$ D projective geometry, the basic invariants are brackets of length n , or in coordinate form, the $n \times n$ determinants formed by the homogeneous coordinates of n vector variables. A bracket polynomial is in normal form if when each term is up to coefficient written in Young tableau form, the entries in each row are increasing, while the entries in each column are non-decreasing [14]. For example for vector variables $\mathbf{v}_1 \prec \mathbf{v}_2 \prec \dots \prec \mathbf{v}_m$, a bracket monomial $[\mathbf{v}_{i_{11}} \mathbf{v}_{i_{12}} \dots \mathbf{v}_{i_{1c}}][\mathbf{v}_{i_{21}} \mathbf{v}_{i_{22}} \dots \mathbf{v}_{i_{2c}}] \dots [\mathbf{v}_{i_{r1}} \mathbf{v}_{i_{r2}} \dots \mathbf{v}_{i_{rc}}]$, where the $\mathbf{v}_{i_{jk}}$ are repetitive selections of the m vector variables, is normal if and only if in

$$\begin{bmatrix} \mathbf{v}_{i_{11}} & \mathbf{v}_{i_{12}} & \dots & \mathbf{v}_{i_{1c}} \\ \mathbf{v}_{i_{21}} & \mathbf{v}_{i_{22}} & \dots & \mathbf{v}_{i_{2c}} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{v}_{i_{r1}} & \mathbf{v}_{i_{r2}} & \dots & \mathbf{v}_{i_{rc}} \end{bmatrix}, \quad (1.1)$$

$\mathbf{v}_{i_{j1}} \prec \mathbf{v}_{i_{j2}} \prec \dots \prec \mathbf{v}_{i_{jc}}$, while $\mathbf{v}_{i_{1k}} \preceq \mathbf{v}_{i_{2k}} \preceq \dots \preceq \mathbf{v}_{i_{rk}}$.

In the advanced invariant theory for 3D orthogonal geometry, each “elementary” advanced invariant is a bracket of length > 1 , whose entries are vector variables representing 3D points. In a bracket monomial, different brackets many have different lengths, and a bracket monomial is in normal form if and only if not only the entries in each row are increasing, the entries in each column are non-decreasing, but all the entries in the tableau after removing the first column, are non-decreasing. For example if (1.1) is normal in this setting, then the sequence $\check{\mathbf{v}}_{i_{11}} \mathbf{v}_{i_{12}} \dots \mathbf{v}_{i_{1c}} \check{\mathbf{v}}_{i_{21}} \mathbf{v}_{i_{22}} \dots \mathbf{v}_{i_{2c}} \dots \check{\mathbf{v}}_{i_{r1}} \mathbf{v}_{i_{r2}} \dots \mathbf{v}_{i_{rc}}$ is non-decreasing, where $\check{\mathbf{v}}_{i_k}$ denotes that $\mathbf{v}_{i_k}$ does not occur in the sequence.

This paper is organized as follows. Section 2 introduces orthogonal geometric invariants by quaternions, Clifford algebra and bracket algebra. Section 3 introduces the main results in [7] on vector-variable polynomials and basics of “advanced bracket algebra”. Section 4 provides the Gröbner base and normal forms of long brackets in the multilinear case. Section 6 extends the results to general case, by means of the square-free vector-variable polynomials introduced in Section 5. Section 7 proposes a normalization algorithm for bracket polynomials.

2. QUATERNIONS, CLIFFORD ALGEBRA, AND BRACKET ALGEBRA

In the vector algebra over $\mathbb{R}^3$, there are three multilinear products among vectors: (i) the inner product of two vectors, (ii) the cross product of two vectors, (iii) the hybrid product of three vectors. None of them can be extended to include more vectors while preserving the associativity.

The quaternionic product, on the other hand, is associative while still multilinear. Let $\bar{\mathbf{q}}$ represent the quaternionic conjugate of quaternion $\mathbf{q}$. Among quaternions, a *vector* $\mathbf{v}$ refers to a pure imaginary quaternion, *i.e.*, $\bar{\mathbf{v}} = -\mathbf{v}$, and a *scalar* $\mathbf{s}$ refers to a real quaternion, *i.e.*, $\bar{\mathbf{s}} = \mathbf{s}$. All vectors span a 3D real inner-product space with metric $\text{diag}(-1, -1, -1)$, denoted by $\mathbb{R}^{-3}$.

We always use juxtaposition of elements to denote their quaternionic product. The *inner product* of two vectors $\mathbf{v}_i, \mathbf{v}_j$ is defined by

$$[\mathbf{v}_i \mathbf{v}_j] := (\mathbf{v}_i \mathbf{v}_j + \mathbf{v}_j \mathbf{v}_i)/2. \quad (2.1)$$

The *cross product* of two vectors $\mathbf{v}_i, \mathbf{v}_j$ is defined by

$$\mathbf{v}_i \times \mathbf{v}_j := (\mathbf{v}_i \mathbf{v}_j - \mathbf{v}_j \mathbf{v}_i)/2. \quad (2.2)$$

The result is a vector, so its inner product with a third vector $\mathbf{v}_k$ is a scalar. Define the *hybrid product* of three vectors $\mathbf{v}_i, \mathbf{v}_j, \mathbf{v}_k$ by

$$[\mathbf{v}_i \mathbf{v}_j \mathbf{v}_k] := (\mathbf{v}_i \mathbf{v}_j \mathbf{v}_k - \mathbf{v}_k \mathbf{v}_j \mathbf{v}_i)/2. \quad (2.3)$$

Then $[\mathbf{v}_i \mathbf{v}_j \mathbf{v}_k] = [(\mathbf{v}_i \times \mathbf{v}_j) \mathbf{v}_k]$.

The vector algebra over $\mathbb{R}^{-3}$ is included in the quaternions. The latter is equipped with a powerful product, the quaternionic product, making it possible to use quaternions to represent 3D orthogonal transformations [1].

The *magnitude* of a quaternion $\mathbf{q}$ is $\sqrt{\bar{\mathbf{q}}\mathbf{q}}$. A quaternion $\mathbf{q}$ is said to be *unit* if $\bar{\mathbf{q}}\mathbf{q} = 1$. Let $\mathbf{q}$ be a unit quaternion, and $\mathbf{v}$ be a vector. The *conjugate adjoint action* of $\mathbf{q}$ on $\mathbf{v}$ is defined by

$$Ad_{\mathbf{q}}(\mathbf{v}) := \bar{\mathbf{q}}\mathbf{v}\mathbf{q}. \quad (2.4)$$

Since $[Ad_{\mathbf{q}}(\mathbf{v}_1) Ad_{\mathbf{q}}(\mathbf{v}_2)] = [\mathbf{v}_1 \mathbf{v}_2]$ for any two vectors $\mathbf{v}_1, \mathbf{v}_2$, $Ad_{\mathbf{q}}$ realizes an orthogonal transformation in $\mathbb{R}^{-3}$. A classical result states that in fact all orthogonal transformations in $\mathbb{R}^{-3}$ are realized in this way, and two different unit quaternions realize the same orthogonal transformation if and only if they differ by sign.

For a quaternion $\mathbf{Q}$, the *bracket* $[\mathbf{Q}]$ is its scalar part:

$$[\mathbf{Q}] := (\mathbf{Q} + \bar{\mathbf{Q}})/2. \quad (2.5)$$

The *axis* of $\mathbf{Q}$ is the vector part of $\mathbf{Q}$:

$$A(\mathbf{Q}) := (\mathbf{Q} - \bar{\mathbf{Q}})/2. \quad (2.6)$$

In particular, $A(\mathbf{v}_1 \mathbf{v}_2) = \mathbf{v}_1 \times \mathbf{v}_2$. We interpret them in geometrical terms below.

For a unit vector $\mathbf{v}_1$, $Ad_{\mathbf{v}_1}$ realizes the reflection with respect to the plane normal to $\mathbf{v}_1$. In general, for unit vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{2k+1}$, $Ad_{\mathbf{v}_1 \mathbf{v}_2 \dots \mathbf{v}_{2k+1}}$ realizes the reflection with respect to the plane normal to axis $A(\mathbf{v}_1 \mathbf{v}_2 \dots \mathbf{v}_{2k+1})$, if the latter is nonzero.

For two unit vectors $\mathbf{v}_1, \mathbf{v}_2$ that are linearly independent, $Ad_{\mathbf{v}_1 \mathbf{v}_2}$ realizes the rotation about the axis $\mathbf{v}_1 \times \mathbf{v}_2$: in the plane spanned by $\mathbf{v}_1, \mathbf{v}_2$, the rotation is from $\mathbf{v}_1$ to the reflection of $\mathbf{v}_1$ with respect to $\mathbf{v}_2$, *i.e.*, the angle of rotation is $\theta = 2\angle(\mathbf{v}_1, \mathbf{v}_2)$. Furthermore, $[\mathbf{v}_1 \mathbf{v}_2] = \cos(\theta/2)$. When we say “rotation $\mathbf{v}_1 \mathbf{v}_2$ ”, we mean the one induced by $Ad_{\mathbf{v}_1 \mathbf{v}_2}$.

In general, for unit vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{2k}$, $Ad_{\mathbf{v}_1 \mathbf{v}_2 \dots \mathbf{v}_{2k}}$ realizes the rotation about the axis $A(\mathbf{v}_1 \mathbf{v}_2 \dots \mathbf{v}_{2k})$, if the latter is nonzero. The rotation is the composition of k rotations $\mathbf{v}_1 \mathbf{v}_2, \mathbf{v}_3 \mathbf{v}_4, \dots, \mathbf{v}_{2k-1} \mathbf{v}_{2k}$. If the angle of rotation is θ , then

$$[\mathbf{v}_1 \dots \mathbf{v}_{2k}] = \cos(\theta/2), \quad |A(\mathbf{v}_1 \dots \mathbf{v}_{2k})| = |\sin(\theta/2)|. \quad (2.7)$$

Let $A(\mathbf{v}_1\mathbf{v}_2\cdots\mathbf{v}_{2k}) \neq 0$. By $[\mathbf{v}_1\mathbf{v}_2\cdots\mathbf{v}_{2k+1}] = [A(\mathbf{v}_1\mathbf{v}_2\cdots\mathbf{v}_{2k})\mathbf{v}_{2k+1}]$, we get

$$[\mathbf{v}_1\mathbf{v}_2\cdots\mathbf{v}_{2k+1}] = \cos \angle(A(\mathbf{v}_1\mathbf{v}_2\cdots\mathbf{v}_{2k}), \mathbf{v}_{2k+1}) \sin(\theta/2), \quad (2.8)$$

where θ is the angle of rotation $\mathbf{v}_1\mathbf{v}_2\cdots\mathbf{v}_{2k}$. In particular when $k = 1$, for linearly independent unit vectors $\mathbf{v}_1, \mathbf{v}_2$, $\sin(\theta/2) = \sin \angle(\mathbf{v}_1, \mathbf{v}_2)$ equals the area of the parallelogram spanned by $\mathbf{v}_1, \mathbf{v}_2$, and $\cos \angle(\mathbf{v}_1 \times \mathbf{v}_2, \mathbf{v}_3)$ equals the height from the end of unit vector $\mathbf{v}_3$ to the plane spanned by $\mathbf{v}_1, \mathbf{v}_2$, so $[\mathbf{v}_1\mathbf{v}_2\mathbf{v}_3]$ equals the volume of the parallelepiped spanned by $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$.

In classical invariant theory, an *algebraic invariant* is a polynomial whose variables are basic invariants. In 3D orthogonal geometry, there are two kinds of basic invariants: $[\mathbf{v}_i\mathbf{v}_j]$ and $[\mathbf{v}_i\mathbf{v}_j\mathbf{v}_k]$ for all vector variables $\mathbf{v}_i, \mathbf{v}_j, \mathbf{v}_k$. Given n vector variables $\mathbf{v}_1, \dots, \mathbf{v}_n$, the brackets $[\mathbf{v}_{j_1}\mathbf{v}_{j_2}\cdots\mathbf{v}_{j_m}]$ for arbitrary $1 < m < \infty$ and arbitrary repetitive selection of elements $\mathbf{v}_{j_1}, \mathbf{v}_{j_2}, \dots, \mathbf{v}_{j_m}$ from the n variables, form an infinite set of advanced algebraic invariants. That each of them is a polynomial of the $[\mathbf{v}_i\mathbf{v}_j]$ and $[\mathbf{v}_i\mathbf{v}_j\mathbf{v}_k]$ is guaranteed by the following *Caianiello expansion formulas* [2], [6]: let $\mathbf{V}_k = \mathbf{v}_{j_1}\mathbf{v}_{j_2}\cdots\mathbf{v}_{j_k}$, then

$$\begin{aligned} [\mathbf{V}_{2l}] &= \sum_{i=2}^{2l} (-1)^i [\mathbf{v}_{j_1}\mathbf{v}_{j_i}][\mathbf{v}_{j_2}\mathbf{v}_{j_3}\cdots\mathbf{v}_{j_i}\cdots\mathbf{v}_{j_{2l}}]; \\ A(\mathbf{V}_{2l-1}) &= \sum_{(2l-2,1) \vdash \mathbf{V}_{2l-1}} [\mathbf{V}_{2l+1(1)}] \mathbf{V}_{2l-1(2)}; \\ A(\mathbf{V}_{2l}) &= \sum_{(2l-2,2) \vdash \mathbf{V}_{2l}} [\mathbf{V}_{2l(1)}] A(\mathbf{V}_{2l(2)}); \\ [\mathbf{V}_{2l+1}] &= \sum_{(2l-2,3) \vdash \mathbf{V}_{2l+1}} [\mathbf{V}_{2l+1(1)}] [\mathbf{V}_{2l+1(2)}], \end{aligned} \quad (2.9)$$

where (i) $(h, m-h) \vdash \mathbf{V}_m$ is a bipartition of the m elements in the sequence $\mathbf{V}_m$ into two subsequences $\mathbf{V}_{m(1)}$ and $\mathbf{V}_{m(2)}$ of length h and $m-h$ respectively; (ii) in $[\mathbf{V}_{m(1)}]$, the product of the h elements in the subsequence is denoted by the same symbol $\mathbf{V}_{m(1)}$; (iii) the summation $\sum_{(h, m-h) \vdash \mathbf{V}_m}$ is over all such bipartitions of $\mathbf{V}_m$, and the sign of permutation of the new sequence $\mathbf{V}_{m(1)}, \mathbf{V}_{m(2)}$ is assumed to be carried by the first factor $[\mathbf{V}_{m(1)}]$ of the addend.

While quaternions are sufficient for describing orthogonal transformations in 3D, they cannot be generalized to higher dimensions directly. In quaternions, the hybrid product $[\mathbf{v}_i\mathbf{v}_j\mathbf{v}_k]$ is a scalar. To make high-dimensional generalization this requirement must be removed, at the same time the property that this element be in the center of the algebra needs to be preserved. If we denote the quaternions by $\mathcal{Q}$, then the above revision leads to a new algebra $\mathcal{Q} \oplus \iota\mathcal{Q}$ of dimension 8, where $\iota := [\mathbf{v}_1\mathbf{v}_2\mathbf{v}_3]$ for three fixed vector variables that are linearly independent. This algebra is the *Clifford algebra* over $\mathbb{R}^{-3}$.

The formal definition of the *Clifford algebra* $\text{Cl}(\mathcal{V}^n)$ over an n -dimensional $\mathbb{K}$ -linear space $\mathcal{V}^n$, where the characteristic of $\mathbb{K}$ is $\neq 2$, is the quotient of the tensor algebra $\bigotimes \mathcal{V}^n$ over the ideal generated by elements of the form $\mathbf{v} \otimes \mathbf{v} - Q(\mathbf{v})$ where Q is a $\mathbb{K}$ -quadratic form. The product induced from the tensor product is called the *Clifford product*, also denoted by juxtaposition of elements [4], [8].

When $\mathcal{V}^n = \mathbb{R}^{-3}$, the quaternionic product of vectors is the image of their Clifford product under the homomorphism induced by mapping ι to a nonzero scalar. In Clifford algebra, ι is not a scalar, but called a *pseudoscalar* because it not

only commutes with everything, but spans a 1D real space containing all hybrid products. The concept quaternionic conjugate is replaced by the *Clifford conjugate*, which is the linear extension of the following operation: for any vectors $\mathbf{v}_{j_1}, \mathbf{v}_{j_2}, \dots, \mathbf{v}_{j_k}$, let $\mathbf{V}_k = \mathbf{v}_{j_1}\mathbf{v}_{j_2}\cdots\mathbf{v}_{j_k}$, then

$$\overline{\mathbf{V}_k} := (-1)^k \mathbf{V}_k^\dagger, \text{ where } \mathbf{V}_k^\dagger := \mathbf{v}_{i_k}\cdots\mathbf{v}_{i_2}\mathbf{v}_{i_1} \text{ is the reversion of } \mathbf{V}_k. \quad (2.10)$$

With the Clifford conjugate, we can define the *magnitude* of $\mathbf{V}_k$, the *conjugate adjoint action* $\text{Ad}_{\mathbf{V}_k}$, the *bracket* $[\mathbf{V}_k]$, the *axis* $A(\mathbf{V}_k)$, together with the concepts not involving conjugate: the inner product $[\mathbf{v}_i\mathbf{v}_j]$, the cross product $\mathbf{v}_i \times \mathbf{v}_j$ and the hybrid product $[\mathbf{v}_i\mathbf{v}_j\mathbf{v}_k]$, just the same as in the case of quaternions. The only difference is that since $[\mathbf{v}_i\mathbf{v}_j\mathbf{v}_k]$ is now a pseudoscalar, while $A(\mathbf{V}_{2l-1})$ remains a vector, $A(\mathbf{V}_{2l})$ is not, but a *pseudovector*. Geometrically, when $A(\mathbf{v}_1\mathbf{v}_2) \neq 0$, it represents the plane spanned by vectors $\mathbf{v}_1, \mathbf{v}_2$, or equivalently, the invariant plane of rotation $\text{Ad}_{\mathbf{v}_1\mathbf{v}_2}$ supporting $\mathbf{v}_1, \mathbf{v}_2$.

We see that unlike the quaternions where there are only two kinds of objects of different dimensions: scalars which are usually called 0-D objects, and vectors which represent 1-D directions and so are usually called 1-D objects, in Clifford algebra $\text{Cl}(\mathbb{R}^{-3})$ there are four kinds of objects of different dimensions. Besides scalars and vectors, there are pseudovectors which represent 2-D directions (planes), and pseudoscalars which represent 3-D orientations (spaces). This is the reason why $\text{Cl}(\mathbb{R}^{-3})$ can be extended to higher dimensions by being capable of discerning objects of different dimensions.

To represent algebraic invariants in 3D orthogonal geometry, using the quaternionic product or the Clifford product in the brackets does not make any difference. The geometric interpretations (2.7), (2.8) and the Caianiello expansion formulas are identical for both products. This justifies the use of juxtaposition of elements to represent both products.

The two kinds of basic invariants $[\mathbf{v}_i\mathbf{v}_j]$ and $[\mathbf{v}_i\mathbf{v}_j\mathbf{v}_k]$ form a commutative ring, called *inner-product bracket algebra*. Formally, given a set of n symbols $\mathcal{M} = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$, two kinds of new symbols can be defined as following: (1) all 2-tuples of symbols selected repetitively from $\mathcal{M}$, by requiring that each 2-tuple be symmetric with respect to its two entries; such a 2-tuple is denoted by $[\mathbf{v}_i\mathbf{v}_j]$. (2) All 3-tuples of symbols selected repetitively from $\mathcal{M}$, by requiring that each 3-tuple is anti-symmetric with respect to its three entries, and in particular, if there are identical entries in a 3-tuple, setting the 3-tuple to be zero; such a 3-tuple is denoted by $[\mathbf{v}_i\mathbf{v}_j\mathbf{v}_k]$.

The two kinds of symbols must satisfy the following *dimension-three constraints*:

IGP : For any five symbols $\mathbf{v}_{i_1}, \dots, \mathbf{v}_{i_5}$,

$$\begin{aligned} & [\mathbf{v}_{i_1}\mathbf{v}_{i_2}][\mathbf{v}_{i_3}\mathbf{v}_{i_4}\mathbf{v}_{i_5}] - [\mathbf{v}_{i_1}\mathbf{v}_{i_3}][\mathbf{v}_{i_2}\mathbf{v}_{i_4}\mathbf{v}_{i_5}] \\ & + [\mathbf{v}_{i_1}\mathbf{v}_{i_4}][\mathbf{v}_{i_2}\mathbf{v}_{i_3}\mathbf{v}_{i_5}] - [\mathbf{v}_{i_1}\mathbf{v}_{i_5}][\mathbf{v}_{i_2}\mathbf{v}_{i_3}\mathbf{v}_{i_4}] = 0. \end{aligned} \quad (2.11)$$

DB : For any six symbols $\mathbf{v}_1, \dots, \mathbf{v}_6$,

$$[\mathbf{v}_1 \mathbf{v}_2 \mathbf{v}_3][\mathbf{v}_4 \mathbf{v}_5 \mathbf{v}_6] = - \begin{vmatrix} [\mathbf{v}_1 \mathbf{v}_4] & [\mathbf{v}_1 \mathbf{v}_5] & [\mathbf{v}_1 \mathbf{v}_6] \\ [\mathbf{v}_2 \mathbf{v}_4] & [\mathbf{v}_2 \mathbf{v}_5] & [\mathbf{v}_2 \mathbf{v}_6] \\ [\mathbf{v}_3 \mathbf{v}_4] & [\mathbf{v}_3 \mathbf{v}_5] & [\mathbf{v}_3 \mathbf{v}_6] \end{vmatrix}. \quad (2.12)$$

The *inner-product bracket algebra* is the commutative ring generated by the above two kinds of symbols, satisfying the symmetry requirements and the dimension-three constraints.

To include brackets of longer length, the concept quaternionic bracket algebra or Clifford bracket algebra needs to be introduced. As explained before, there is no need to distinguish between the two concepts in the setting of 3D orthogonal geometry, so we simply call it *bracket algebra*. To distinguish from the concept of the same name arising from Grassmann-Cayley algebra [14], we call that in [14] *classical bracket algebra*.

Formally, besides the above 2-tuples and 3-tuples, a hierarchy of infinitely many new symbols can be defined: for any length $l > 3$, there are all l -tuples of symbols selected repetitively from $\mathcal{M}$, with the requirement that the first and the last equalities in Caianiello expansion (2.9) are satisfied; such an l -tuple is denoted by $[\mathbf{v}_{j_1} \mathbf{v}_{j_1} \cdots \mathbf{v}_{j_l}]$. By setting $[1] = 1$ (0-tuple) and $[\mathbf{v}_i] = 0$ (1-tuple) for all i , we get a full hierarchy of new symbols marked by brackets, with arbitrary length $l \geq 0$. The new symbols together with their specific requirements, form a commutative ring called the *bracket algebra* over 3D inner-product space. This is the bottom-up approach to defining bracket algebra. The concept quaternionic product or Clifford product is not needed.

3. VECTOR-VARIABLE POLYNOMIALS AND BRACKET POLYNOMIALS

In this paper, we use (1) bold-faced digital numbers and bold-faced lower-case letters to denote vector variables, *e.g.*, $\mathbf{v}, 1$; (2) bold-faced upper-case letters to denote monic vector-variable monomials, *e.g.*, $\mathbf{A}, \mathbf{X}$; (3) Roman-styled lower-case letters to denote polynomials, *e.g.*, f, g ; (4) Greek letters to denote $\mathbb{K}$ -coefficients, *e.g.*, λ, μ .

Although the background is real orthogonal geometry, the algebraic manipulations under investigation are independent of the real field. In fact, only the following coefficients occur in computing: $\pm 2^k$ for $k \in \mathbb{Z}$. We set the base field $\mathbb{K}$ to be of characteristic $\neq 2$.

Now start from quaternions. Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ be symbols. What properties determine that the multilinear associative product among the symbols is the quaternionic one, and that these symbols represent vectors of a 3D real inner-product space with metric $\text{diag}(-1, -1, -1)$? [7] gives a rather simple answer.

The inner product of two vectors $\mathbf{v}_i, \mathbf{v}_j$ is a scalar, so it commutes with a third vector $\mathbf{v}_k$: $[\mathbf{v}_i \mathbf{v}_j] \mathbf{v}_k = \mathbf{v}_k [\mathbf{v}_i \mathbf{v}_j]$. For three vectors $\mathbf{v}_i, \mathbf{v}_j, \mathbf{v}_k$, since $[\mathbf{v}_i \mathbf{v}_j \mathbf{v}_k]$ is a scalar, for a fourth vector $\mathbf{v}_l$, the commutativity $[\mathbf{v}_i \mathbf{v}_j \mathbf{v}_k] \mathbf{v}_l = \mathbf{v}_l [\mathbf{v}_i \mathbf{v}_j \mathbf{v}_k]$ holds. The two commutativities are all that characterize the equal-

ity properties of the quaternionic product, besides the multilinearity and associativity.

THEOREM 1. [7] *Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ be $n > 2$ symbols. Define the product among them, denoted by juxtaposition of elements, as the $\mathbb{K}$ -tensor product modulo the two-sided ideal generated by the following tensors:*

$$\begin{aligned} \mathbf{V}2 : & \quad \mathbf{v}_i \otimes \mathbf{v}_i \otimes \mathbf{v}_j - \mathbf{v}_j \otimes \mathbf{v}_i \otimes \mathbf{v}_i; \\ \mathbf{V}3 : & \quad (\mathbf{v}_i \otimes \mathbf{v}_j + \mathbf{v}_j \otimes \mathbf{v}_i) \otimes \mathbf{v}_k - \mathbf{v}_k \otimes (\mathbf{v}_i \otimes \mathbf{v}_j + \mathbf{v}_j \otimes \mathbf{v}_i); \\ \mathbf{V}4 : & \quad (\mathbf{v}_i \otimes \mathbf{v}_j \otimes \mathbf{v}_k - \mathbf{v}_k \otimes \mathbf{v}_j \otimes \mathbf{v}_i) \otimes \mathbf{v}_l \\ & \quad - \mathbf{v}_l \otimes (\mathbf{v}_i \otimes \mathbf{v}_j \otimes \mathbf{v}_k - \mathbf{v}_k \otimes \mathbf{v}_j \otimes \mathbf{v}_i), \end{aligned} \quad (3.1)$$

for any $i \neq j \neq k \neq l$ in $1, 2, \dots, n$. Denote by $\mathcal{Q}$ or $\mathcal{Q}[[\mathbf{v}_1, \dots, \mathbf{v}_n]]$ the $\mathbb{K}$ -algebra defined by the above product and generated by the $\mathbf{v}_i$. Denote by $\mathcal{I}[[\mathbf{v}_1, \dots, \mathbf{v}_n]]$ the above ideal, and call it the *syzygy ideal* of $\mathcal{Q}$.

Denote

$$\begin{aligned} \mathbb{K}_2 &:= \mathbb{K}(\{\mathbf{v}_i \otimes \mathbf{v}_i, \mathbf{v}_i \otimes \mathbf{v}_j + \mathbf{v}_j \otimes \mathbf{v}_i, |i \neq j\}), \\ \mathbb{K}_3 &:= \mathbb{K}_2(\{\mathbf{v}_i \otimes \mathbf{v}_j \otimes \mathbf{v}_k - \mathbf{v}_k \otimes \mathbf{v}_j \otimes \mathbf{v}_i | i \neq j \neq k\}). \end{aligned} \quad (3.2)$$

(1) Let $\iota := [\mathbf{v}_1 \mathbf{v}_2 \mathbf{v}_3]$. Then $\iota \neq 0$, and $\mathbb{K}_3 = \mathbb{K}_2(\iota)$.

(2) The $\mathbf{v}_i$ and $\iota \mathbf{v}_i \times \mathbf{v}_j$ of the $\mathbb{K}$ -algebra $\mathcal{Q}$ span a 3D $\mathbb{K}_2$ -space $\mathcal{V}^3$. For any $k \geq 1$, $A(\mathbf{v}_{j_1} \cdots \mathbf{v}_{j_{2k+1}})$ and $\iota A(\mathbf{v}_{j_1} \cdots \mathbf{v}_{j_{2k}})$ are both in $\mathcal{V}^3$.

(3) The defined product is the Clifford product of the $\mathbb{K}_2$ -Clifford algebra over $\mathcal{V}^3$.

(4) If $\mathbb{K} = \mathbb{K}_3 = \mathbb{R}$, and the inner product of real space $\mathcal{V}^3$ induced from (2.1) is definite, then the defined product is the quaternionic product.

(5) Let $\mathbf{V}_k = \mathbf{v}_{j_1} \mathbf{v}_{j_2} \cdots \mathbf{v}_{j_k}$. Then the following identities hold modulo $\mathcal{I}[[\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]]$:

$$\mathbf{v}_i [\mathbf{V}_k] = [\mathbf{V}_k] \mathbf{v}_i, \quad [\mathbf{v}_i \mathbf{V}_k] = [\mathbf{V}_k \mathbf{v}_i]. \quad (3.3)$$

The requirements in (4) distinguishing the quaternionic product from the Clifford product cannot be represented by equalities. So for symbolic manipulations of equalities, the quaternionic product and the Clifford product cannot be distinguished. The $\mathbb{K}$ -algebra $\mathcal{Q}$ is called the *3D vector-variable polynomial ring* generated by vector variables $\mathbf{v}_i$, and the product in it is called the *vector-variable product*. It is neither the quaternionic product nor the Clifford product, but a more basic one.

All the terminologies introduced earlier on quaternions and Clifford algebra are valid for $\mathcal{Q}$. Besides, there are some additional terminologies for $\mathcal{Q}$. Let $\mathbf{v}_1 \prec \mathbf{v}_2 \prec \dots \prec \mathbf{v}_n$ be vector variables. A *monic monomial* of vector variables refers to the product of a repetitive permutation of some of the vector variables. For a monic monomial $\mathbf{v}_{i_1} \mathbf{v}_{i_2} \cdots \mathbf{v}_{i_k}$, the *leading variable* refers to $\mathbf{v}_{i_1}$, and the *trailing variable* refers to $\mathbf{v}_{i_k}$. The monomial is said to be *non-descending* if $\mathbf{v}_{i_1} \preceq \mathbf{v}_{i_2} \preceq \dots \preceq \mathbf{v}_{i_k}$, and is said to be *ascending* if $\mathbf{v}_{i_1} \prec \mathbf{v}_{i_2} \prec \dots \prec \mathbf{v}_{i_k}$. The *degree*, or *length*, of the monomial is k . The *lexicographic ordering* among monomials is always assumed.

A *polynomial* of vector variables is a $\mathbb{K}$ -linear combination of monic monomials. The *leading term* of a polynomial f is the term of highest order, denoted by $\text{lt}(f)$. The *degree* of a polynomial is that of its leading term. The leading terms of all elements in a subset $\mathcal{S}$ of polynomials are denoted by $\text{lt}(\mathcal{S})$. When specifying the field $\mathbb{K}_2$ or $\mathbb{K}_3$, we can get the corresponding concepts *quaternionic polynomial* and *Clifford polynomial*.

Fix a multiset of vector variables $\mathcal{M}$ composed of $m \geq 3$ symbols $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$. Let n be the number of different elements in $\mathcal{M}$, where $3 \leq n \leq m$. In the $\mathbb{K}$ -tensor algebra $\otimes(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n)$ generated by the n symbols taken as vectors, a *tensor monomial* is up to coefficient the tensor product of finitely many such vectors. The $\mathbb{K}$ -*tensor algebra* over $\mathcal{M}$, denoted by $\otimes[\mathcal{M}]$, is the $\mathbb{K}$ -subspace of $\otimes(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n)$ spanned by tensor monomials whose vector variables by counting multiplicity are in $\mathcal{M}$, equipped with the tensor product that is undefined if the result is no longer in $\otimes[\mathcal{M}]$.

When the product among the elements in $\mathcal{M}$ is the vector-variable product, we have the corresponding concept of *3D vector-variable polynomial ring over multiset $\mathcal{M}$* , denoted by $\mathcal{Q}[\mathcal{M}]$. Each element in $\mathcal{Q}[\mathcal{M}]$ is a polynomial whose multiset of vector variables in each term is a submultiset of $\mathcal{M}$. The syzygy ideal $\mathcal{I}[\mathcal{M}]$ of $\mathcal{Q}[\mathcal{M}]$ is still defined by (3.1).

The concepts of *Gröbner base* and *normal form* are defined in $\mathcal{Q}[\mathcal{M}]$ just as in $\otimes(\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n)$ [9]. For two monomials h_1, h_2 in vector variables, h_1 is said to be *reduced* with respect to h_2 , if h_2 is not a *factor* of h_1 , or h_1 is not a *multiplier* of h_2 , i.e., there do not exist monomials l, r , including elements of $\mathbb{K}$, such that $h_1 = lh_2r$. For two polynomials f and g , f is said to be *reduced* with respect to g , if the leading term of f is reduced with respect to that of g . The term “*non-reduced*” means the opposite.

Let $\{f_1, f_2, \dots, f_k\}$ be a set of vector-variable polynomials. Another set of vector-variable polynomials $\{g_1, g_2, \dots, g_l\}$ is said to be a *reduced Gröbner base* of the ideal $\mathcal{I} := \langle f_1, f_2, \dots, f_k \rangle$ generated by the f_i in $\mathcal{Q}[\mathcal{M}]$, if (1) $\langle g_1, \dots, g_l \rangle = \mathcal{I}$, (2) the leading term of any element in $\mathcal{I}$ is a multiplier of the leading term of some g_i , (3) the g_i are pairwise reduced with respect to each other.

The *reduction* of a polynomial f with respect to a reduced Gröbner base $g_1, g_2, \dots, g_l$ is the repetitive procedure of dividing the highest-ordered non-reduced term L of f by some g_i whose leading term is a factor of L , then updating f by replacing L with its remainder, until all terms of f are reduced. The result is called the *normal form* of f with respect to the Gröbner base. Two polynomials are equal if and only if they have identical normal forms.

In [7], two theorems are established for the Gröbner base and normal forms of 3D vector-variable polynomials, one for the multilinear case where each element in multiset $\mathcal{M}$ has multiplicity 1, the other for the general case $\mathcal{Q}[\mathbf{v}_1, \dots, \mathbf{v}_n]$.

THEOREM 2. [7] *Let $\mathcal{I}[\mathbf{v}_1, \dots, \mathbf{v}_n]$ be the syzygy ideal of the multilinear polynomial ring $\mathcal{Q}[\mathbf{v}_1, \dots, \mathbf{v}_n]$ in n different vector variables $\mathbf{v}_1 \prec \mathbf{v}_2 \prec \dots \prec \mathbf{v}_n$.*

(1) [Gröbner base] *The following are a reduced Gröbner base of $\mathcal{I}[\mathbf{v}_1, \dots, \mathbf{v}_n]$: for all $1 \leq i_1 < i_2 < \dots < i_j \leq n$,*

$$\text{G3: } [\mathbf{v}_{i_3} \mathbf{v}_{i_2} \mathbf{v}_{i_1}] - [\mathbf{v}_{i_1} \mathbf{v}_{i_3} \mathbf{v}_{i_2}], \text{ and } [\mathbf{v}_{i_3} \mathbf{v}_{i_1} \mathbf{v}_{i_2}] - [\mathbf{v}_{i_2} \mathbf{v}_{i_3} \mathbf{v}_{i_1}];$$

$$\text{Gj: } [\mathbf{v}_{i_3} \mathbf{v}_{i_2} \mathbf{v}_{i_4} \mathbf{v}_{i_5} \cdots \mathbf{v}_{i_j} \mathbf{v}_{i_1}] - [\mathbf{v}_{i_2} \mathbf{v}_{i_4} \mathbf{v}_{i_5} \cdots \mathbf{v}_{i_j} \mathbf{v}_{i_1} \mathbf{v}_{i_3}], \text{ for all } j > 3.$$

(2) [Normal form] *In a normal form, every term is up to coefficient of the form $\mathbf{V}_{Y_1} \mathbf{v}_{z_1} \mathbf{V}_{Y_2} \mathbf{v}_{z_2} \cdots \mathbf{V}_{Y_k} \mathbf{v}_{z_k}$ or $\mathbf{V}_{Y_1} \mathbf{v}_{z_1} \cdots \mathbf{V}_{Y_k} \mathbf{v}_{z_k} \mathbf{V}_{Y_{k+1}}$, where*

- (i) $k \geq 0$,
- (ii) $\mathbf{v}_{z_1} \mathbf{v}_{z_2} \cdots \mathbf{v}_{z_k}$ is ascending,
- (iii) every $\mathbf{V}_{Y_i}$ is an ascending monomial of length ≥ 1 ,
- (iv) $\mathbf{V}_{Y_1} \mathbf{V}_{Y_2} \cdots \mathbf{V}_{Y_k}$ (or $\mathbf{V}_{Y_1} \mathbf{V}_{Y_2} \cdots \mathbf{V}_{Y_{k+1}}$ if $\mathbf{V}_{Y_{k+1}}$ occurs) is ascending,
- (v) for every $i \leq k$, if $\mathbf{v}_{t_i}$ is the trailing variable of monomial $\mathbf{V}_{Y_i}$, then $\mathbf{v}_{z_i} \prec \mathbf{v}_{t_i}$.

THEOREM 3. [7] *Let $\mathcal{I}[\mathbf{v}_1, \dots, \mathbf{v}_n]$ be the syzygy ideal of the polynomial ring $\mathcal{Q}[\mathbf{v}_1, \dots, \mathbf{v}_n]$ in n different vector variables $\mathbf{v}_1 \prec \mathbf{v}_2 \prec \dots \prec \mathbf{v}_n$.*

(1) [Gröbner base] *The following are a reduced Gröbner base of $\mathcal{I}[\mathbf{v}_1, \dots, \mathbf{v}_n]$:*

$$\text{G3, Gj: for all } 3 < j < \infty, \text{ and all } 1 \leq i_1 < i_2 < i_3 < i_4 \leq \dots \leq i_{j-1} < i_j \leq n;$$

$$\text{EG2: for all } i_1 < i_2,$$

$$\mathbf{v}_{i_2} \mathbf{v}_{i_2} \mathbf{v}_{i_1} - \mathbf{v}_{i_1} \mathbf{v}_{i_2} \mathbf{v}_{i_2}, \quad \mathbf{v}_{i_2} \mathbf{v}_{i_1} \mathbf{v}_{i_1} - \mathbf{v}_{i_1} \mathbf{v}_{i_1} \mathbf{v}_{i_2};$$

$$\text{EGj: } [\mathbf{v}_{i_3} \mathbf{v}_{i_2} \mathbf{v}_{i_3} \mathbf{v}_{i_4} \cdots \mathbf{v}_{i_j} \mathbf{v}_{i_1}] - [\mathbf{v}_{i_2} \mathbf{v}_{i_3} \mathbf{v}_{i_4} \cdots \mathbf{v}_{i_j} \mathbf{v}_{i_1} \mathbf{v}_{i_3}], \text{ for all } 2 < j < \infty, \text{ and all } 1 \leq i_1 < i_2 < i_3 \leq i_4 \leq \dots \leq i_{j-1} < i_j \leq n.$$

(2) [Normal form] *In a normal form, every term is up to coefficient of the form $\mathbf{V}_{Y_1} \mathbf{v}_{h_1} \mathbf{v}_{z_1} \mathbf{V}_{Y_2} \mathbf{v}_{h_2} \mathbf{v}_{z_2} \cdots \mathbf{V}_{Y_k} \mathbf{v}_{h_k} \mathbf{v}_{z_k}$ or $\mathbf{V}_{Y_1} \mathbf{v}_{h_1} \mathbf{v}_{z_1} \cdots \mathbf{V}_{Y_k} \mathbf{v}_{h_k} \mathbf{v}_{z_k} \mathbf{V}_{Y_{k+1}}$, where*

- (i) $k \geq 0$,
- (ii) $\mathbf{v}_{z_1} \mathbf{v}_{z_2} \cdots \mathbf{v}_{z_k}$ is non-descending,
- (iii) $\mathbf{v}_{h_1} \mathbf{v}_{h_2} \cdots \mathbf{v}_{h_k}$ is non-descending,
- (iv) every $\mathbf{V}_{Y_i}$ is a non-descending monomial of length ≥ 0 ,
- (v) $\mathbf{V}_{Y_1} \mathbf{v}_{h_1} \mathbf{V}_{Y_2} \mathbf{v}_{h_2} \cdots \mathbf{V}_{Y_k} \mathbf{v}_{h_k}$ (or $\mathbf{V}_{Y_1} \mathbf{v}_{h_1} \cdots \mathbf{V}_{Y_k} \mathbf{v}_{h_k} \mathbf{V}_{Y_{k+1}}$ if $\mathbf{V}_{Y_{k+1}}$ occurs) is non-descending,
- (vi) for every $i \leq k$, $\mathbf{v}_{h_i} \succ \mathbf{v}_{z_i}$,
- (vii) for every $i \leq k$, if the length of $\mathbf{V}_{Y_i}$ is nonzero, let $\mathbf{v}_{t_i}$ be the trailing variable of $\mathbf{V}_{Y_i}$, then $\mathbf{v}_{t_i} \prec \mathbf{v}_{h_i}$.

For a general multiset $\mathcal{M}$ in which the different vector variables are $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$, the Gröbner base of the syzygy ideal $\mathcal{I}[\mathcal{M}]$ is the restriction of the Gröbner base of $\mathcal{I}[\mathbf{v}_1, \dots, \mathbf{v}_n]$ to $\mathcal{Q}[\mathcal{M}]$, denoted by $\mathcal{G}[\mathcal{M}]$. In $\mathcal{Q}[\mathcal{M}]$, a polynomial is said to be $\mathcal{I}$ -*normal* if its leading term is reduced with respect to the Gröbner base. The procedure of deriving the normal form of a polynomial is called $\mathcal{I}$ -*reduction*.

Now that any vector-variable polynomial has a normal form by $\mathcal{I}$ -reduction, so does a bracket polynomial when every

bracket is expanded into two terms by definition. The result is complicated.

Consider the following simple example: for a single bracket

$$2[\mathbf{v}_1 \mathbf{v}_2 \cdots \mathbf{v}_m] = \mathbf{v}_1 \mathbf{v}_2 \cdots \mathbf{v}_m + (-1)^m \mathbf{v}_m \cdots \mathbf{v}_2 \mathbf{v}_1,$$

the $\mathcal{I}$ -reduction goes as follows: for $0 \leq j < m$, if we define $\mathbf{V}_{m-j} = \mathbf{v}_{j+1} \mathbf{v}_{j+2} \cdots \mathbf{v}_m$, then when $m \geq 3$,

$$\begin{aligned} & \mathbf{V}_m + (-1)^m \mathbf{V}_m^\dagger \\ \stackrel{\mathcal{I}}{=} & \mathbf{v}_1 \mathbf{V}_{m-1} + \mathbf{V}_{m-1} \mathbf{v}_1 - \mathbf{v}_1 (\mathbf{V}_{m-1} + (-1)^{m-1} \mathbf{V}_{m-1}^\dagger) \\ \stackrel{\mathcal{I}}{=} & \mathbf{v}_1 \mathbf{V}_{m-1} + \mathbf{V}_{m-1} \mathbf{v}_1 - \mathbf{v}_1 (\mathbf{v}_2 \mathbf{V}_{m-2} + \mathbf{V}_{m-2} \mathbf{v}_2) \\ & + \mathbf{v}_1 \mathbf{v}_2 (\mathbf{V}_{m-2} + (-1)^{m-2} \mathbf{V}_{m-2}^\dagger) \\ = & \mathbf{v}_2 \mathbf{V}_{m-2} \mathbf{v}_1 - \mathbf{v}_1 \mathbf{V}_{m-2} \mathbf{v}_2 \\ & + \mathbf{v}_1 \mathbf{v}_2 (\mathbf{V}_{m-2} + (-1)^{m-2} \mathbf{V}_{m-2}^\dagger). \end{aligned} \quad (3.4)$$

From this recursive formula, we get for $m \geq 1$,

$$\begin{aligned} 2[\mathbf{v}_1 \mathbf{v}_2 \cdots \mathbf{v}_m] & \stackrel{\mathcal{I}}{=} \frac{1 + (-1)^m}{2} \mathbf{v}_1 \mathbf{v}_2 \cdots \mathbf{v}_m \\ & + \sum_{i=1}^{m-1} (-1)^{i+1} (\mathbf{v}_1 \cdots \mathbf{v}_i \cdots \mathbf{v}_m) \mathbf{v}_i. \end{aligned} \quad (3.5)$$

So the normal form of the simplest bracket $[\mathbf{v}_1 \mathbf{v}_2 \cdots \mathbf{v}_m]$ is composed of up to m terms. What is worse is that only when the terms are summed up can they represent a single algebraic invariant (the bracket), while missing a single term destroys the invariance of the whole expression.

From the appearance, the bracket symbol only hides half of a binomial. There are much more behind this appearance. By definition, for a sequence $\mathbf{A}$ of $a > 1$ vector variables, its bracket is $[\mathbf{A}] = 2^{-1} \mathbf{A} + (-1)^a 2^{-1} \mathbf{A}^\dagger$. Monomial $\mathbf{A}$ is called the *representative* of bracket $[\mathbf{A}]$. Later on, when we write $[\mathbf{A}]$, we always assume that monomial $\mathbf{A}$ is the representative of the bracket.

The definition of a bracket endows the symbol with the reversion symmetry (or equivalently, the conjugate symmetry) up to sign: $[\mathbf{v}_{i_1} \cdots \mathbf{v}_{i_a}] = (-1)^a [\mathbf{v}_{i_a} \cdots \mathbf{v}_{i_1}]$. By (3.3), the bracket symbol also has shift symmetry. So up to sign a bracket of a vector variables has the symmetry group D_{2a} (dihedral group).

The above analysis is only for a single bracket. For bracket polynomials, there are a lot of polynomial identities, or *syzygies*, among them. These complexities justify the separation of bracket algebra from vector-variable polynomial ring in symbolic manipulations of algebraic invariants. Finding the Gröbner base of the syzygies and then characterizing the normal forms of bracket polynomials are the main goal of this paper.

The following are some terminology on brackets. The *representative* of a bracket polynomial is the vector-variable polynomial whose terms are each the product of the coefficient with the representatives of the bracket factors in the same term. The representative of a bracket polynomial is allowed to contain brackets. For example, $[\mathbf{v}_1 \mathbf{v}_2 [\mathbf{v}_3 \mathbf{v}_4]]$ is taken as bracket binomial $2^{-1} [\mathbf{v}_1 \mathbf{v}_2 (\mathbf{v}_3 \mathbf{v}_4 + \mathbf{v}_4 \mathbf{v}_3)]$; its representative is the content $\mathbf{v}_1 \mathbf{v}_2 [\mathbf{v}_3 \mathbf{v}_4]$ within the outer bracket.

The *lexicographic ordering* of bracket polynomials is that of their representatives. The *leading variable* of a bracket refers to that of its representative. The *leading term* of a bracket polynomial is always under the lexicographic ordering. For example, if $\mathbf{v}_1 \prec \mathbf{v}_2$, then $[\mathbf{v}_1 \mathbf{v}_2] \prec [\mathbf{v}_2 \mathbf{v}_1]$, and $[\mathbf{v}_1 \mathbf{v}_2][\mathbf{v}_2 \mathbf{v}_1] \prec [\mathbf{v}_2 \mathbf{v}_1][\mathbf{v}_1 \mathbf{v}_2]$.

The *leader* (or *expanded leading term*) of a bracket polynomial, refers to the leading term of the bracket polynomial when taken as a vector-variable one, *i.e.*, the corresponding vector-variable polynomial obtained from expanding each bracket into two terms by definition. For example, the leader of bracket $[\mathbf{A}]$ refers to the one of higher order between $2^{-1} \mathbf{A}$ and $(-1)^a 2^{-1} \mathbf{A}^\dagger$.

Among all the bracket polynomials that are equal to the same bracket polynomial, there are two that have strong features: the first is the one whose representative is the lowest, the second is the one whose leader is the lowest. The second is unique but the first is not. To make the first unique we introduce the following concept.

In $\mathcal{Q}[\mathcal{M}]$, where the number of elements in multiset $\mathcal{M}$ is m , a *uni-bracket monomial* refers to a single bracket of length m . A *uni-bracket polynomial* is a $\mathbb{K}$ -linear combination of uni-bracket monomials. All uni-bracket polynomials form a $\mathbb{K}$ -linear space, denoted by $[\mathcal{Q}][\mathcal{M}]$. The $\mathbb{K}$ -linear space of the representatives of elements in $[\mathcal{Q}][\mathcal{M}]$ is just the $\mathbb{K}$ -linear space of degree- m vector-variable polynomials, denoted by $\mathcal{Q}_m[\mathcal{M}]$. Obviously, $[\mathcal{Q}][\mathcal{M}]$ is a linear subspace of $\mathcal{Q}_m[\mathcal{M}]$.

When taken as a vector-variable polynomial, a uni-bracket is a binomial. In appearance, a uni-bracket is a monomial. To distinguish between the two understandings, we need a device to get rid of the bracket symbol and extract the representative of the uni-bracket. This can be done by taking $[\mathcal{Q}][\mathcal{M}]$ as the quotient of $\mathcal{Q}_m[\mathcal{M}]$ modulo the ideal

$$\mathcal{J}[\mathcal{M}] := \mathcal{I}[\mathcal{M}] + [\mathcal{I}][\mathcal{M}], \quad (3.6)$$

where $[\mathcal{I}][\mathcal{M}]$ is composed of the vector parts of degree- m polynomials, *i.e.*, the $\mathbb{K}$ -linear span of elements of the form

$$\mathbf{R} : \mathbf{v}_{i_1} \mathbf{v}_{i_2} \cdots \mathbf{v}_{i_m} - (-1)^m \mathbf{v}_{i_m} \cdots \mathbf{v}_{i_2} \mathbf{v}_{i_1}, \quad (3.7)$$

for all permutations of the m elements in $\mathcal{M}$.

The modulo- $[\mathcal{I}][\mathcal{M}]$ operation identifies a uni-bracket with its representative, or equivalently, identifies any degree- m vector-variable polynomial with the uni-bracket polynomial it serves as the representative. This operation retains the bracket symbols of all the brackets of length $< m$, while removing the bracket symbols from all brackets of length m .

4. GRÖBNER BASE AND NORMAL FORM FOR MULTILINEAR UNI-BRACKET POLYNOMIALS

From this section on, we use bold-faced digital numbers to denote vector variables, and use bold-faced capital letters to denote monic monomials of vector variables.

In this section, the multiset $\mathcal{M}$ is composed of $m \geq 3$ different vector variables, and the modulo- $[\mathcal{I}][\mathcal{M}]$ operation is always assumed. Then $[\mathcal{Q}][\mathcal{M}]$ and $\mathcal{Q}_m[\mathcal{M}]$ are identical, and

a uni-bracket no longer has the outer bracket symbol. Ideal $[\mathcal{I}][\mathcal{M}]$ is called the *uni-bracket removal ideal* in $\mathcal{Q}[\mathcal{M}]$, and ideal $\mathcal{J}[\mathcal{M}]$ is called the *syzygy ideal* of $[\mathcal{Q}][\mathcal{M}]$ in $\otimes[\mathcal{M}]$. Below we compute the Gröbner base of $\mathcal{J}[\mathcal{M}]$ and characterize the normal forms of uni-bracket polynomials.

The following are elements of $\mathcal{J}[\mathcal{M}]$: for all monomials $\mathbf{A}$ to $\mathbf{F}$ such that $\mathbf{A1B}, \mathbf{1C}, \mathbf{1D2}, \mathbf{1E2F}$ are of length m and $\mathbf{E}$ has length $e > 0$,

$$\begin{aligned} S_1 : & \quad \mathbf{A1B} - \mathbf{1BA}, \\ S_1^N : & \quad \text{the } S_1 \text{ in which } \mathbf{A1B} \text{ is } \mathcal{I}\text{-normal}, \\ R_1 : & \quad 2(\mathbf{1[C]}) = \mathbf{1C} - (-1)^m \mathbf{1C}^\dagger, \\ R_{12} : & \quad 2(\mathbf{1[D2]}) = \mathbf{1D2} - (-1)^m \mathbf{12D}^\dagger, \\ R_{12}[*] : & \quad 2(\mathbf{1[E2[F]]}) = \mathbf{1E2[F]} - (-1)^e \mathbf{12E}^\dagger[\mathbf{F}]. \end{aligned} \quad (4.1)$$

Since the reduced Gröbner base of $\mathcal{I}[\mathcal{M}]$ is $\mathcal{G}[\mathcal{M}]$, we only need to consider the elements of type R in $\mathcal{J}[\mathcal{M}]$, as they span $[\mathcal{I}][\mathcal{M}]$. By

$$\begin{aligned} & \mathbf{A1B} - (-1)^m \mathbf{B}^\dagger \mathbf{1A}^\dagger \\ = & (\mathbf{A1B} - \mathbf{1BA}) + (\mathbf{1BA} - (-1)^m \mathbf{1A}^\dagger \mathbf{B}^\dagger) \\ & - (-1)^m (\mathbf{B}^\dagger \mathbf{1A}^\dagger - \mathbf{1A}^\dagger \mathbf{B}^\dagger), \end{aligned} \quad (4.2)$$

we get

LEMMA 4. R is a subset of the ideal $\langle S_1, R_1 \rangle$. For any type-R element f but not of type R_1 , $\text{lt}(f) \in \text{lt}(S_1)$.

LEMMA 5. S_1 is a subset of $\langle S_1^N, R_1 \rangle + \mathcal{I}[\mathcal{M}]$. For any type- S_1 element f but not of type S_1^N , $\text{lt}(f) \in \text{lt}(\mathcal{I}[\mathcal{M}])$.

Proof. Consider a general type- S_1 element $f = \mathbf{A1B} - \mathbf{1BA}$, where $\mathbf{A}$ is not empty. If $\mathbf{A1B}$ is $\mathcal{I}$ -normal, then $f \in S_1^N$. If $\mathcal{I}$ -reductions are carried out to $\mathbf{A}, \mathbf{B}$, say $\mathbf{A} = \mathbf{A}^N + \mathcal{I}$ and $\mathbf{B} = \mathbf{B}^N + \mathcal{I}$, where $\mathbf{A}^N, \mathbf{B}^N$ are both $\mathcal{I}$ -normal, then $f = \mathbf{A}^N \mathbf{1B}^N - \mathbf{1B}^N \mathbf{A}^N + \mathcal{I}$, and $\mathbf{A}^N \mathbf{1B}^N$ is the new leading term. So by $\mathcal{I}$ -reductions we can assume that both $\mathbf{A}$ and $\mathbf{B}$ are $\mathcal{I}$ -normal.

Assume that $\mathbf{A1B}$ is not $\mathcal{I}$ -normal. Further assume that any type- S_1 element $g \prec f$ is in $\langle S_1^N, R_1 \rangle + \mathcal{I}[\mathcal{M}]$. We prove the conclusion for f by reduction on the order. There are three possibilities to apply Gröbner base elements of $\mathcal{I}[\mathcal{M}]$ to make reduction to $\mathbf{A1B}$: (i) G3 at the end of $\mathbf{A1}$, (ii) G_i for $i > 3$ at the end of $\mathbf{A1}$, (iii) G3 on 1 and two variables from $\mathbf{A}, \mathbf{B}$ respectively.

Case (i). Let $\mathbf{A1B} = \mathbf{Cuv1B}$, where $\mathbf{u} \succ \mathbf{v} \succ \mathbf{1}$. Then

$$\begin{aligned} f & \stackrel{G_3}{=} -\mathbf{Cvu1B} + \mathbf{C1(uv + vu)B} - \mathbf{1BCuv} \\ & \stackrel{\text{induction}}{=} -\mathbf{1BC(vu + uv)} + \mathbf{1(uv + vu)BC} \\ & \stackrel{\mathcal{I}}{=} 0. \end{aligned}$$

Case (ii). Let $\mathbf{A1B} = \mathbf{CuvD1B}$, where $\mathbf{1} \prec \mathbf{v} \prec \mathbf{u} \prec$ all variables of $\mathbf{D}$. Let the length of $\mathbf{D}$ be $d > 0$. Let the lengths

of $\mathbf{B}, \mathbf{C}$ be b, c respectively. Then $b + c = m - d - 3$.

$$\begin{aligned} f & \stackrel{G_i}{=} \mathbf{C(vD1 + (-1)^d \mathbf{1D}^\dagger \mathbf{v})uB} - (-1)^d \mathbf{Cu1D}^\dagger \mathbf{vB} \\ & \quad - \mathbf{1BCuvD} \\ & \stackrel{\text{induction}}{=} \mathbf{1\{(uBC - BCu)vD + (-1)^d \mathbf{D}^\dagger \mathbf{v}(uBC - BCu)\}} \\ & \stackrel{R_1}{=} (-1)^d \mathbf{1D}^\dagger \mathbf{v}\{(-1)^{m-d}(\mathbf{C}^\dagger \mathbf{B}^\dagger \mathbf{u} - \mathbf{uC}^\dagger \mathbf{B}^\dagger) \\ & \quad + (\mathbf{uBC} - \mathbf{BCu})\} \\ & = (-1)^d \mathbf{1D}^\dagger \mathbf{v}\{-((-1)^{b+c} \mathbf{C}^\dagger \mathbf{B}^\dagger + \mathbf{BC})\mathbf{u} \\ & \quad + \mathbf{u}((-1)^{b+c} \mathbf{C}^\dagger \mathbf{B}^\dagger + \mathbf{BC})\} \\ & \stackrel{\mathcal{I}}{=} 0. \end{aligned}$$

Case (iii). Let $\mathbf{A1B} = \mathbf{Cu1vD}$, where $\mathbf{u} \succ \mathbf{v} \succ \mathbf{1}$. Let the lengths of $\mathbf{C}, \mathbf{D}$ be c, d respectively. Then $c + d = m - 3$.

$$\begin{aligned} f & \stackrel{G_3}{=} -\mathbf{C1uvD} + \mathbf{Cv(u1 + 1u)D} - \mathbf{1vDCu} \\ & \stackrel{\text{induction}}{=} \mathbf{1(-uvDC + DCvu + uDCv - vDCu)} \\ & \stackrel{R_1}{=} \mathbf{1\{-((-1)^m \mathbf{C}^\dagger \mathbf{D}^\dagger - \mathbf{DC})vu} \\ & \quad + \mathbf{v}((-1)^m \mathbf{C}^\dagger \mathbf{D}^\dagger - \mathbf{DC})\mathbf{u}\}} \\ & \stackrel{\mathcal{I}}{=} 0. \end{aligned}$$

□

LEMMA 6. R_1 is a subset of the ideal $\langle R_{12} \rangle + \mathcal{I}[\mathcal{M}]$. For any type- R_1 element f but not of type R_{12} , $\text{lt}(f) \in \text{lt}(\mathcal{I}[\mathcal{M}])$.

Proof. Let $f = 2(\mathbf{1[E2F]})$ be a general element of type R_1 . Then $f \stackrel{\mathcal{I}}{=} 2(\mathbf{1[FE2]})$, and the latter is in R_{12} . □

THEOREM 7. Let $\mathcal{M} = \{\mathbf{1}, \mathbf{2}, \dots, \mathbf{m}\}$ be $m > 2$ different symbols, where $\mathbf{1} \prec \mathbf{2} \prec \dots \prec \mathbf{m}$, and let $\mathcal{Q}[\mathcal{M}]$ be the vector-variable polynomial ring over $\mathcal{M}$. Let $[\mathcal{Q}][\mathcal{M}]$ be the space of uni-bracket polynomials, and let $\mathcal{J}[\mathcal{M}]$ be its syzygy ideal in $\otimes[\mathcal{M}]$.

(1) [Gröbner base] The following are a reduced Gröbner base of $\mathcal{J}[\mathcal{M}]$:

$\mathcal{G}[\mathcal{M}]$: G_i for all $3 \leq i \leq m$;

S_1^N : $\mathbf{A1B} - \mathbf{1BA}$, where $\mathbf{A}$ is an ascending sequence of length $a > 0$, $\mathbf{B}$ is of length $m - a - 1 \geq 0$, and $\mathbf{A1B}$ is $\mathcal{I}$ -normal;

$R_{12}^N[j]$: for all $1 \leq j \leq \frac{m-1}{2}$, $\mathbf{1[A2][Y_2z_2][Y_3z_3] \cdots [Y_jz_j]}$, where (i) $\mathbf{A}$ is an ascending sequence of length $a > 0$; (ii) when $j = 1$, then $R_{12}^N[1] = \mathbf{1[A2]}$; (iii) each $\mathbf{Y}_i$ is a non-empty ascending sequence, and each $\mathbf{z}_i$ is a variable such that $\mathbf{z}_i \mathbf{Y}_i$ is ascending; (iv) $\mathbf{1A2Y_2z_2 \cdots Y_jz_j}$ is $\mathcal{I}$ -normal and length- m .

(2) [Normal form] In a normal form, every term is $\mathcal{I}$ -normal, and is up to coefficient of one of the following forms:

(I) $\mathbf{12C}$, where $\mathbf{C}$ is $\mathcal{I}$ -normal;

(II) $\mathbf{1A2Y_2z_2Y_3z_3 \cdots Y_{k-1}z_{k-1}Y_k}$, where $k \geq 2$, $\mathbf{A}$ and the $\mathbf{Y}_i$ are each a non-empty ascending sequence, and each $\mathbf{z}_i \prec \mathbf{t}_i$, the latter being the trailing variable of $\mathbf{Y}_i$;

(III) $\mathbf{1A2Y_2Z_2Y_3Z_3 \cdots Y_kZ_k}$, where $k \geq 2$, $\mathbf{A}$ and the $\mathbf{Y}_i, \mathbf{z}_i$ are as in (II), and for some $2 \leq i \leq k$, if $\mathbf{l}_i$ is the leading variable of $\mathbf{Y}_i$, then $\mathbf{l}_i \prec \mathbf{z}_i$.

Proof. There are several steps.

Step 1. We need to prove that R_{12} is a subset of the ideal $\langle R_{12}^N \rangle + \mathcal{I}[\mathcal{M}]$. Once this is done, then since the leader of any element of R_{12}^N is $\mathcal{I}$ -normal and cannot be cancelled by the leader of any other element of R_{12}^N , the G_i and $R_{12}^N[j]$ form a reduced Gröbner base of $\langle R_{12} \rangle + \mathcal{I}[\mathcal{M}]$. By this and the previous three lemmas, we get conclusion (1) of the theorem.

Once conclusion (1) holds, then any $\mathcal{I}$ -normal monomial of length m with leading variable $\mathbf{1}$ is the representative of a uni-bracket in normal form if and only if it is not the leader of an R_{12}^N -typed element. Conclusion (2) follows.

Step 2. The idea of proving the statement in Step 1 is to use $\mathcal{G}[\mathcal{M}]$ to decrease the order of the leader of every element of type R_{12} or $R_{12}[*]$, at the same time keep the reduction result to be within the $\mathbb{K}$ -linear space spanned by elements of type R_{12} or $R_{12}[*]$. Then ultimately all the leaders of these elements become $\mathcal{I}$ -normal.

We start with the $\mathcal{I}$ -reduction on the leading term of a general R_{12} -typed element $f = \mathbf{1A2} - (-1)^m \mathbf{12A}^\dagger$, where the length of $\mathbf{A}$ is $m - 2$. If by $\mathcal{I}$ -reduction, $\mathbf{A} = \mathbf{A}^N + \mathcal{I}$, then $f = \mathbf{1A}^N \mathbf{2} - (-1)^m \mathbf{12A}^N + \mathcal{I}$. So we can assume that $\mathbf{A}$ is $\mathcal{I}$ -normal.

If $\mathbf{A2}$ is $\mathcal{I}$ -normal, then f is just $R_{12}^N[1]$. When $\mathbf{A2}$ is not $\mathcal{I}$ -normal, if $\mathbf{A2}$ is non-reduced with respect to G_3 , let $\mathbf{A2} = \mathbf{Buv2}$, where $\mathbf{u} \succ \mathbf{v} \succ \mathbf{2}$, then

$$\mathbf{1Buv2} = -\mathbf{1Bvu2} + 2\{\mathbf{1B2[uv]}\}. \quad (4.3)$$

The result consists of the leading terms of one R_{12} -typed element and one $R_{12}[*]$ -typed element. The leaders of both terms are lower than f .

If $\mathbf{A2}$ is non-reduced with respect to G_i for some $i > 3$, let $\mathbf{A2} = \mathbf{BuvC2}$, where $\mathbf{u} \succ \mathbf{v} \succ \mathbf{2}$, and $\mathbf{uC}$ is ascending, and the length of $\mathbf{C}$ is $c > 0$. Then

$$\begin{aligned} \mathbf{1BuvC2} &\stackrel{G_i}{=} \mathbf{1B(vC2 + (-1)^c \mathbf{2C}^\dagger \mathbf{v})u} - (-1)^c \mathbf{1Bu2C}^\dagger \mathbf{v} \\ &= \frac{\mathbf{1BvC2u}}{\mathbf{1B2uvC}} + \mathbf{1B2((-1)^c \mathbf{C}^\dagger \mathbf{vu} + \mathbf{uvC})} \\ &\quad - \frac{\mathbf{1B2uvC}}{\mathbf{1Bu2vC}} + \mathbf{1Bu2(-(-1)^c \mathbf{C}^\dagger \mathbf{v} + \mathbf{vC})} \\ &\stackrel{\mathcal{I}}{=} 2\{\mathbf{1B2[uvC]}\} + 2\{\mathbf{1Bu2[vC]}\} - \mathbf{1BvCu2}, \end{aligned} \quad (4.4)$$

The result consists of the leading terms of two R_{12} -typed elements and one $R_{12}[*]$ -typed element. The leaders of the three terms are lower than f .

By (4.3) and (4.4), a monomial that is non-reduced with respect to a G_i for some $i \geq 3$ must contain a subsequence of the form $\mathbf{uDv}$, where (i) the length of $\mathbf{D}$ is $d > 0$; (ii) $\mathbf{u} \succ \mathbf{v}$; (iii) if $\mathbf{l_D}$ is the leading variable of $\mathbf{D}$, then $\mathbf{u} \succ \mathbf{l_D}$; (iv) if $\mathbf{D}$ contains more than one variable, then $\mathbf{l_D} \succ \mathbf{v}$. (4.3) and (4.4) can be written in the following unified form:

$$\mathbf{uDv} \stackrel{G(d+2)}{=} 2(\mathbf{uv[D]}) + 2(\mathbf{v[uD]}) - \mathbf{Duv}. \quad (4.5)$$

It is called the *fundamental $\mathcal{I}$ -reduction formula*.

Step 3. Consider $\mathcal{I}$ -reductions on the leader of a general $R_{12}[*]$ -typed element $f = \mathbf{1A2[B]} - (-1)^a \mathbf{12A}^\dagger[\mathbf{B}]$, where the length of $\mathbf{B}$ is $b > 0$.

Since $[\mathbf{B}] = (-1)^b [\mathbf{B}^\dagger]$, henceforth we assume that in any $R_{12}[*]$ -typed element to be normalized, the leading variable of any bracket has higher order than the trailing variable of the bracket. Then the leader of the bracket is always its representative.

In this step, we consider $\mathcal{I}$ -reduction to the representative $\mathbf{B}$ of $[\mathbf{B}]$. Let $[\mathbf{B}] = [\mathbf{CuDvE}]$. Substituting (4.5) into it, we get

$$[\mathbf{CuDvE}] \stackrel{\mathcal{I}}{=} 2[\mathbf{CuvE}][\mathbf{D}] + 2[\mathbf{CvE}][\mathbf{uD}] - [\mathbf{CDuvE}]. \quad (4.6)$$

Step 4. Consider $\mathcal{I}$ -reductions on the leader $\mathbf{1A2B}$ of $f = \mathbf{1A2[B]} - (-1)^a \mathbf{12A}^\dagger[\mathbf{B}]$ involving both the tail part of $\mathbf{A}$ and the head part of $\mathbf{B}$, where the leading variable of $\mathbf{B}$ is assumed to be higher than the trailing variable.

As $\mathbf{2}$ is lower than any element of $\mathbf{A}, \mathbf{B}$, the only possible reduction is by G_3 . Let $\mathbf{1A2B} = \mathbf{1Ca2bD}$, where $\mathbf{C}$ may be empty but $\mathbf{D}$ is not. Assume $\mathbf{a} \succ \mathbf{b} \succ \mathbf{t_D}$, where $\mathbf{t_D}$ is the trailing variable of $\mathbf{D}$. Let the length of $\mathbf{D}$ be d . It is easy to prove that applying G_3 to $\mathbf{a2b}$ in vector-variable binomial $\mathbf{1Ca2[bD]}$ is equivalent to the following *absorption of bracket*:

$$\begin{aligned} \mathbf{1Ca2[bD]} &\stackrel{\mathcal{I}}{=} \mathbf{1C[bD]a2} \\ &= 2^{-1}(\mathbf{1CbDa2}) + (-1)^d 2^{-1}(\mathbf{1CD}^\dagger \mathbf{ba2}). \end{aligned} \quad (4.7)$$

Each term in the result is a leading term of an R_{12} -typed element lower than f .

Step 5. In Step 3, we have seen that a single bracket after $\mathcal{I}$ -reduction, may be split into two brackets. The split can continue and we gradually get expressions of the form

$$R_{12}[j] : \mathbf{1E2[F_2][F_3] \cdots [F_j]} - (-1)^e \mathbf{12E}^\dagger[\mathbf{F_2}][\mathbf{F_3}] \cdots [\mathbf{F_j}], \quad (4.8)$$

where the length of $\mathbf{E}$ is $e > 0$, and the length of $\mathbf{1E2F_2F_3 \cdots F_j}$ is m . $R_{12}[j]$ is a $\mathbb{K}$ -linear combination of elements of type $R_{12}[*]$ if all but one bracket are each expanded into two terms.

Consider $\mathcal{I}$ -reductions of $R_{12}[j]$ involving more than two bracket factors, and $\mathcal{I}$ -reductions involving $\mathbf{1E2}$ and more than one bracket factor. Since $\mathbf{2}$ is lower than all elements of $\mathbf{E}$ and the $\mathbf{F_i}$, G_3 is the only possible Gröbner base element that may apply to $\mathbf{2}$ and its neighbors on both sides simultaneously. G_3 can involve only $[\mathbf{F_2}]$ among the brackets.

In $[\mathbf{F_2}][\mathbf{F_3}] \cdots [\mathbf{F_j}]$, only G_i where $i > 3$ can involve more than two brackets. However, since G_i is of the form $\mathbf{uDv}$ where $\mathbf{D}$ is ascending, if the product of the leaders of three brackets is non-reduced with respect to some G_i , then the middle bracket must be composed of a subsequence of $\mathbf{D}$ of length ≥ 2 , contradicting with the assumption that the leading variable in the middle bracket be higher than the trailing variable.

So each $\mathcal{I}$ -reduction of $R_{12}[j]$ by a single Gi where $i \geq 3$, can involve at most two bracket factors, or the $\mathbf{1E2}$ and one bracket factor.

Step 6. Consider $\mathcal{I}$ -reductions on $[\mathbf{F}_1][\mathbf{F}_2]$, where the leading variable in each bracket is higher than the trailing variable. If the leading variable $\mathbf{l}_{\mathbf{F}_1}$ of $\mathbf{F}_1$ is higher than the leading variable $\mathbf{l}_{\mathbf{F}_2}$ of $\mathbf{F}_2$, then an $\mathcal{I}$ -reduction commuting the two brackets reduces the order of their product. Below we always assume $\mathbf{l}_{\mathbf{F}_1} \prec \mathbf{l}_{\mathbf{F}_2}$.

For $G3$, there are two possibilities to involve both $\mathbf{F}_1, \mathbf{F}_2$ in the leader $\mathbf{F}_1\mathbf{F}_2$: two variables at the end of $\mathbf{F}_1$ and the third at the beginning of $\mathbf{F}_2$, or one variable at the end of $\mathbf{F}_1$ and the other two at the beginning of $\mathbf{F}_2$. The latter is impossible because $\mathbf{l}_{\mathbf{F}_1} \prec \mathbf{l}_{\mathbf{F}_2}$. For Gi where $i > 3$, there are also two possibilities: two variables at the end of $\mathbf{F}_1$ and the rest at the beginning of $\mathbf{F}_2$, or one variable at the end of $\mathbf{F}_1$ and the rest at the beginning of $\mathbf{F}_2$. The latter is also impossible due to $\mathbf{l}_{\mathbf{F}_1} \prec \mathbf{l}_{\mathbf{F}_2}$.

Case $G3$. Let $[\mathbf{F}_1][\mathbf{F}_2] = [\mathbf{Buv}][\mathbf{wCd}]$, where $\mathbf{u} \succ \mathbf{v}$, and $\mathbf{u} \succ \mathbf{w} \succ \mathbf{d}$. Let the length of $\mathbf{C}$ be $c \geq 0$, and let the leading variable of $\mathbf{Bu}$ be $\mathbf{l}$. Then $\mathbf{w} \succ \mathbf{l} \succ \mathbf{v}$. Applying $G3$ to $\mathbf{u}\mathbf{v}\mathbf{w}$ is equivalent to the following *absorption of the second bracket*:

$$\begin{aligned} & [\mathbf{Buv}][\mathbf{wCd}] \\ \stackrel{\mathcal{I}}{=} & [\mathbf{B}[\mathbf{wCd}]\mathbf{uv}] \\ = & 2^{-1}[\mathbf{BwCd}]\mathbf{uv} + (-1)^{c-1}[\mathbf{BdC}^\dagger\mathbf{wuv}]. \end{aligned} \quad (4.9)$$

The leader of each bracket monomial in the result has lower order than the leader of $[\mathbf{F}_1][\mathbf{F}_2]$.

Case Gi . Let $[\mathbf{F}_1][\mathbf{F}_2] = [\mathbf{Buv}][\mathbf{aDwC}]$, where $\mathbf{aD}$ is ascending, and $\mathbf{a} \succ \mathbf{u} \succ \mathbf{v} \succ \mathbf{w}$. Let the lengths of $\mathbf{B}, \mathbf{C}, \mathbf{D}$ be b, c, d respectively. Let the leading variable of $\mathbf{Bu}$ be $\mathbf{l}$, and let the trailing variable of $\mathbf{wC}$ be $\mathbf{t}$. Then $\mathbf{a} \succ \mathbf{l} \succ \mathbf{v}$ and $\mathbf{a} \succ \mathbf{t}$. Applying $G(d+4)$ to $\mathbf{u}\mathbf{v}\mathbf{aDw}$, we get

$$\begin{aligned} & 4[\mathbf{Buv}][\mathbf{aDwC}] \\ = & \mathbf{Bu}\mathbf{aDwC} + (-1)^{c+d}\mathbf{Bu}\mathbf{vC}^\dagger\mathbf{wD}^\dagger\mathbf{a} \\ & + (-1)^b\mathbf{v}\mathbf{uB}^\dagger\mathbf{aDwC} + (-1)^{b+c+d}\mathbf{v}\mathbf{uB}^\dagger\mathbf{C}^\dagger\mathbf{wD}^\dagger\mathbf{a} \\ \stackrel{\mathcal{I}}{=} & (\mathbf{v}\mathbf{aDw} - (-1)^d\mathbf{wD}^\dagger\mathbf{av})\mathbf{BuC} + (-1)^d\mathbf{Bu}\mathbf{wD}^\dagger\mathbf{avC} \\ & + (-1)^d((-1)^c\mathbf{vC}^\dagger\mathbf{w} + \mathbf{wCv})\mathbf{BuD}^\dagger\mathbf{a} \\ & - (-1)^d\mathbf{Bu}\mathbf{wCvD}^\dagger\mathbf{a} \\ & + (-1)^b\mathbf{v}(\mathbf{aDwC} + (-1)^{c+d}\mathbf{C}^\dagger\mathbf{wD}^\dagger\mathbf{a})\mathbf{uB}^\dagger \\ = & \mathbf{v}\{\mathbf{aDw}(\mathbf{BuC} - (-1)^{b+c}\mathbf{C}^\dagger\mathbf{uB}^\dagger) \\ & + (-1)^b\mathbf{aDw}(\mathbf{C} + (-1)^c\mathbf{C}^\dagger)\mathbf{uB}^\dagger \\ & + (-1)^{c+d}\mathbf{C}^\dagger\mathbf{w}(\mathbf{BuD}^\dagger\mathbf{a} + (-1)^{b+d}\mathbf{aDuB}^\dagger) \\ & - (-1)^{b+c}\mathbf{C}^\dagger\mathbf{w}(\mathbf{aD} - (-1)^d\mathbf{D}^\dagger\mathbf{a})\mathbf{uB}^\dagger\} \\ & + (-1)^d\mathbf{w}\{-\mathbf{D}^\dagger\mathbf{av}(\mathbf{BuC} - (-1)^{b+c}\mathbf{C}^\dagger\mathbf{uB}^\dagger) \\ & - (-1)^{b+c}\mathbf{D}^\dagger\mathbf{avC}^\dagger\mathbf{uB}^\dagger - (-1)^{b+d}\mathbf{CvaDuB}^\dagger \\ & + \mathbf{Cv}(\mathbf{BuD}^\dagger\mathbf{a} + (-1)^{b+d}\mathbf{aDuB}^\dagger)\} \\ & + (-1)^d\mathbf{Bu}\mathbf{w}(\mathbf{D}^\dagger\mathbf{avC} - \mathbf{CvD}^\dagger\mathbf{a}) \\ \stackrel{\mathcal{I}}{=} & 4([\mathbf{v}\mathbf{aDw}][\mathbf{BuC}] + (-1)^d[\mathbf{wCv}][\mathbf{BuD}^\dagger\mathbf{a}]) \\ & + (-1)^b\mathbf{v}(\mathbf{aDC} + (-1)^{c+d}\mathbf{D}^\dagger\mathbf{aC}^\dagger)\mathbf{wuB}^\dagger \\ & - (-1)^b\mathbf{w}(\mathbf{CvaD} + (-1)^{c+d}\mathbf{D}^\dagger\mathbf{avC}^\dagger)\mathbf{uB}^\dagger \\ & + (-1)^d\mathbf{Bu}\mathbf{w}(\mathbf{D}^\dagger\mathbf{avC} - \mathbf{CvD}^\dagger\mathbf{a}) \end{aligned}$$

$$\begin{aligned} \stackrel{\mathcal{I}}{=} & 4([\mathbf{v}\mathbf{aDw}][\mathbf{BuC}] + (-1)^d[\mathbf{wCv}][\mathbf{BuD}^\dagger\mathbf{a}]) \\ & + (-1)^b2\{\mathbf{v}(\mathbf{aD}[\mathbf{C}] - (-1)^c[\mathbf{aD}]\mathbf{C}^\dagger)\mathbf{wuB}^\dagger\} \\ & - 4[\mathbf{Bu}\mathbf{w}][\mathbf{CvaD}] + \mathbf{Bu}\mathbf{w}\{\mathbf{C}(\mathbf{aD} - (-1)^d\mathbf{D}^\dagger\mathbf{a}) \\ & \quad + (-1)^d\mathbf{D}^\dagger\mathbf{a}(\mathbf{C} + (-1)^c\mathbf{C}^\dagger)\}\mathbf{v} \\ \stackrel{\mathcal{I}}{=} & 4\{[\mathbf{v}\mathbf{aDw}][\mathbf{BuC}] + (-1)^d[\mathbf{wCv}][\mathbf{BuD}^\dagger\mathbf{a}] \\ & \quad - [\mathbf{Bu}\mathbf{w}][\mathbf{CvaD}] + [\mathbf{Bu}\mathbf{w}(\mathbf{C}[\mathbf{aD}] + (-1)^d\mathbf{D}^\dagger\mathbf{a}[\mathbf{C}])\mathbf{v}]\} \\ \stackrel{\mathcal{I}}{=} & 4\{[\mathbf{v}\mathbf{aDw}][\mathbf{BuC}] + (-1)^d[\mathbf{wCv}][\mathbf{BuD}^\dagger\mathbf{a}] \\ & \quad - [\mathbf{Bu}\mathbf{w}][\mathbf{CvaD}] + [\mathbf{Bu}\mathbf{wCv}][\mathbf{aD}] \\ & \quad + (-1)^d[\mathbf{Bu}\mathbf{wD}^\dagger\mathbf{av}][\mathbf{C}]\}. \end{aligned} \quad (4.10)$$

The leader of each bracket monomial in the result has lower order than the leader of $[\mathbf{F}_1][\mathbf{F}_2]$.

Step 7. Consider a bracket of the form $h = [\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1\mathbf{a}_2\mathbf{B}_2\mathbf{c}_2 \cdots \mathbf{a}_k\mathbf{B}_k\mathbf{c}_k]$, where (1) $k > 1$, (2) $\mathbf{a}_i \succ \mathbf{c}_i$ for every i , (3) $\mathbf{a}_i \succ \mathbf{c}_j$ for all $1 \leq j \leq k$. Let the length of $\mathbf{B}_i$ be b_i .

When $k = 2$,

$$\begin{aligned} & [\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1\mathbf{a}_2\mathbf{B}_2\mathbf{c}_2] - 2[\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1][\mathbf{a}_2\mathbf{B}_2\mathbf{c}_2] \\ = & 2^{-1}\{(-1)^{b_1+b_2}\mathbf{c}_2\mathbf{B}_2^\dagger\mathbf{a}_2\mathbf{c}_1\mathbf{B}_1^\dagger\mathbf{a}_1 - (-1)^{b_2}\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1\mathbf{c}_2\mathbf{B}_2^\dagger\mathbf{a}_2 \\ & - (-1)^{b_1}\mathbf{c}_1\mathbf{B}_1^\dagger\mathbf{a}_1\mathbf{a}_2\mathbf{B}_2\mathbf{c}_2 - (-1)^{b_1+b_2}\mathbf{c}_1\mathbf{B}_1^\dagger\mathbf{a}_1\mathbf{c}_2\mathbf{B}_2^\dagger\mathbf{a}_2\} \\ \stackrel{\mathcal{I}}{=} & -(-1)^{b_2}[\mathbf{c}_2\mathbf{B}_2^\dagger\mathbf{a}_2\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1]. \end{aligned}$$

The leader in the result has lower order than h .

For $k > 2$,

$$\begin{aligned} & [\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1 \cdots \mathbf{a}_k\mathbf{B}_k\mathbf{c}_k] \\ & - 2[\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1 \cdots \mathbf{a}_{k-1}\mathbf{B}_{k-1}\mathbf{c}_{k-1}][\mathbf{a}_k\mathbf{B}_k\mathbf{c}_k] \\ \stackrel{\mathcal{I}}{=} & -(-1)^{b_k}[\mathbf{c}_k\mathbf{B}_k^\dagger\mathbf{a}_k(\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1 \cdots \mathbf{a}_{k-1}\mathbf{B}_{k-1}\mathbf{c}_{k-1})], \end{aligned}$$

and for $[\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1 \cdots \mathbf{a}_{k-1}\mathbf{B}_{k-1}\mathbf{c}_{k-1}]$, the split into $[\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1 \cdots \mathbf{a}_{k-2}\mathbf{B}_{k-2}\mathbf{c}_{k-2}][\mathbf{a}_{k-1}\mathbf{B}_{k-1}\mathbf{c}_{k-1}]$ can continue. In the end, we get

$$[\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1 \cdots \mathbf{a}_k\mathbf{B}_k\mathbf{c}_k] \stackrel{\mathcal{I}}{=} 2^{k-1}[\mathbf{a}_1\mathbf{B}_1\mathbf{c}_1] \cdots [\mathbf{a}_k\mathbf{B}_k\mathbf{c}_k] + g, \quad (4.11)$$

where g is a bracket polynomial whose leader has lower order than h .

(4.11) can be used to split a long bracket whose representative is $\mathcal{I}$ -normal. It can also be used in the converse direction, to concatenate short brackets into a long one.

Step 8. So far we have proved that for any R_{12} -typed element $\mathbf{1}[\mathbf{E2}]$ or any $R_{12}[j]$ -typed element $\mathbf{1}[\mathbf{A2}][\mathbf{F2}][\mathbf{F3}] \cdots [\mathbf{Fj}]$, as long as the leader is not $\mathcal{I}$ -normal, $\mathcal{I}$ -reductions can always be carried out to change $\mathbf{1E2}$ or $\mathbf{1A2}[\mathbf{F2}][\mathbf{F3}] \cdots [\mathbf{Fj}]$ into the following form:

$$\begin{aligned} \mathbf{1}T = & \sum_{\alpha} \lambda_{\alpha} \mathbf{1E}_{\alpha} \mathbf{2} + \sum_{\beta} \mu_{\beta} \mathbf{1A}_{\beta} \mathbf{2} [\mathbf{F}_{\beta_2}][\mathbf{F}_{\beta_3}] \cdots [\mathbf{F}_{\beta_j}] \\ & + \sum_{\gamma} \tau_{\gamma} \mathbf{12} [\mathbf{D}_{\gamma_1}][\mathbf{D}_{\gamma_2}] \cdots [\mathbf{D}_{\gamma_k}], \end{aligned} \quad (4.12)$$

where the leading variable in each bracket is higher than the trailing variable, the $\mathbf{1E}_{\alpha} \mathbf{2}$ and $\mathbf{1A}_{\beta} \mathbf{2} \mathbf{F}_{\beta_2} \mathbf{F}_{\beta_3} \cdots \mathbf{F}_{\beta_j}$ are all $\mathcal{I}$ -normal.

Since any $\mathcal{I}$ -normal form is of type $\mathbf{Y}_1\mathbf{z}_1 \cdots \mathbf{Y}_k\mathbf{z}_k$ or $\mathbf{Y}_1\mathbf{z}_1 \cdots \mathbf{Y}_k\mathbf{z}_k \mathbf{Y}_{k+1}$, it must be that

- (i) $\mathbf{1E}_\alpha \mathbf{2} = \mathbf{Y}_1 \mathbf{z}_1$, i.e., $\mathbf{E}_\alpha = \mathbf{34} \cdots \mathbf{m}$.
(ii) $\mathbf{1A}_\beta \mathbf{2F}_{\beta_2} \mathbf{F}_{\beta_3} \cdots \mathbf{F}_{\beta_j} = \mathbf{Y}_1 \mathbf{z}_1 \cdots \mathbf{Y}_j \mathbf{z}_j$, and

$$\begin{cases} \mathbf{1A}_\beta \mathbf{2} &= \mathbf{Y}_1 \mathbf{z}_1, \\ \mathbf{F}_{\beta_2} &= \mathbf{Y}_2 \mathbf{z}_2, \\ \cdots &= \cdots \\ \mathbf{F}_{\beta_j} &= \mathbf{Y}_j \mathbf{z}_j. \end{cases}$$

(i) is obvious. In (ii), the trailing variable of each bracket must be some $\mathbf{z}_i$. If an $\mathbf{F}_{\beta_i}$ is $\mathbf{Y}_h \mathbf{z}_h \cdots \mathbf{Y}_{h+p} \mathbf{z}_{h+p}$ for some $p > 0$, for $h \leq s \leq h+p$, let the leading variable of $\mathbf{Y}_s$ be $\mathbf{l}_s$, then $\mathbf{l}_h \succ \mathbf{z}_{h+p} \succ \mathbf{z}_{h+p-1} \succ \cdots \succ \mathbf{z}_h$. By (4.11), $[\mathbf{F}_{\beta_i}]$ is split into $2^p [\mathbf{Y}_h \mathbf{z}_h] \cdots [\mathbf{Y}_{h+p} \mathbf{z}_{h+p}]$ plus some bracket monomials of lower leader. Then $\mathcal{I}$ -reductions continue to the terms involving such bracket monomials. Ultimately each bracket is of the form $[\mathbf{Y}_i \mathbf{z}_i]$.

In (4.12), $\mathbf{1E}_\alpha \mathbf{2} = (-1)^m \mathbf{1}(\mathbf{E}_\alpha \mathbf{2})^\dagger$ by $R_{12}^N[1]$, $\mathbf{1A}_\beta \mathbf{2} [\mathbf{F}_{\beta_2}] \cdots [\mathbf{F}_{\beta_j}] = (-1)^m \mathbf{1}(\mathbf{A}_\beta \mathbf{2} [\mathbf{F}_{\beta_2}] \cdots [\mathbf{F}_{\beta_j}])^\dagger$ by $R_{12}^N[j]$, and $\mathbf{12}[\mathbf{D}_{\gamma_1}] \cdots [\mathbf{D}_{\gamma_k}] = (-1)^m \mathbf{1}(\mathbf{2}[\mathbf{D}_{\gamma_1}] \cdots [\mathbf{D}_{\gamma_k}])^\dagger$ by $\mathcal{I}[\mathcal{M}]$. So $\mathbf{1T} - (-1)^m \mathbf{1T}^\dagger$ is reduced to zero by $\mathcal{I}[\mathcal{M}]$ and R_{12}^N . $\square$

5. SQUARE-FREE VECTOR-VARIABLE POLYNOMIAL RING

When $\mathcal{M}$ is a general multiset of vector variables, a *square* in $\mathcal{Q}[\mathcal{M}]$ refers to the product of a vector with itself. Denote $\mathbf{v}_i^2 := \mathbf{v}_i \mathbf{v}_i$. It commutes with everything in $\mathcal{Q}[\mathcal{M}]$.

PROPOSITION 8. *In $\mathcal{Q}[\mathcal{M}]$, let $f = g\mathbf{v}_i^2 h$ be a multiplier of $\mathbf{v}_i^2$. If f is not $\mathcal{I}$ -normal, then by doing $\mathcal{I}$ -reduction to g, h , together with rearranging the position of $\mathbf{v}_i^2$ in each term, f can become $\mathcal{I}$ -normal.*

Proof. Suppose g, h are $\mathcal{I}$ -normal. There are three cases for f to be non-reduced with respect to the Gröbner base $\mathcal{G}[\mathcal{M}]$:

(1) If f contains as a factor the leader of Gk for $k > 3$, or EGj for $j > 1$ involving both of $\mathbf{v}_i^2$, then $\mathbf{v}_i^2$ is preserved by the reduction with the Gröbner base element.

(2) If $g\mathbf{v}_i$ is non-reduced, then switch the element of $\mathcal{G}[\mathcal{M}]$ with respect to which $g\mathbf{v}_i$ is non-reduced:

Case EG2: Let $g\mathbf{v}_i = \mathbf{A}\mathbf{u}\mathbf{v}_i \mathbf{v}_i$ or $\mathbf{A}\mathbf{u}\mathbf{v}_i$, where $\mathbf{u} \succ \mathbf{v}_i$. Then $\mathbf{A}\mathbf{u}\mathbf{v}_i \mathbf{v}_i^2 h = \mathbf{A}\mathbf{v}_i^2 \mathbf{u}\mathbf{v}_i h$ or $\mathbf{A}\mathbf{u}\mathbf{v}_i^2 h = \mathbf{A}\mathbf{v}_i^2 \mathbf{u} h$.

Case G3: Let $g\mathbf{v}_i = \mathbf{A}\mathbf{u}\mathbf{w}\mathbf{v}_i$ where $\mathbf{u} \succ \mathbf{w}$ and $\mathbf{u} \succ \mathbf{v}_i$. Then $\mathbf{A}\mathbf{u}\mathbf{w}\mathbf{v}_i^2 h = \mathbf{A}\mathbf{v}_i^2 \mathbf{u}\mathbf{w} h$.

Case Gk for $k > 3$ or EGj for $j > 2$: Let $g\mathbf{v}_i = \mathbf{C}\mathbf{u}\mathbf{w}\mathbf{D}\mathbf{v}_i$ where $\mathbf{u} \succ \mathbf{w} \succ \mathbf{v}_i$. Then $\mathbf{C}\mathbf{u}\mathbf{w}\mathbf{D}\mathbf{v}_i^2 h = \mathbf{C}\mathbf{v}_i^2 \mathbf{u}\mathbf{w}\mathbf{D} h$.

(3) If $\mathbf{v}_i h$ is non-reduced, then switch the element of $\mathcal{G}[\mathcal{M}]$ with respect to which $\mathbf{v}_i h$ is non-reduced:

Case EG2: Let $\mathbf{v}_i h = \mathbf{v}_i \mathbf{v}_i \mathbf{z}\mathbf{B}$ or $\mathbf{v}_i \mathbf{z}\mathbf{z}\mathbf{B}$, where $\mathbf{v}_i \succ \mathbf{z}$. Then $g\mathbf{v}_i^2 \mathbf{v}_i \mathbf{z}\mathbf{B} = g\mathbf{v}_i \mathbf{z}\mathbf{v}_i^2 \mathbf{B}$ or $g\mathbf{v}_i^2 \mathbf{z}\mathbf{z}\mathbf{B} = g\mathbf{z}\mathbf{z}\mathbf{v}_i^2 \mathbf{B}$.

Case G3: Let $\mathbf{v}_i h = \mathbf{v}_i \mathbf{y}\mathbf{z}\mathbf{B}$ where $\mathbf{v}_i \succ \mathbf{y}$ and $\mathbf{v}_i \succ \mathbf{z}$. Then $g\mathbf{v}_i^2 \mathbf{y}\mathbf{z}\mathbf{B} = g\mathbf{y}\mathbf{z}\mathbf{v}_i^2 \mathbf{B}$.

Case Gk for $k > 3$ or EGj for $j > 2$: Let $\mathbf{v}_i h = \mathbf{v}_i \mathbf{y}\mathbf{C}\mathbf{z}\mathbf{D}$ where $\mathbf{v}_i \succ \mathbf{y} \succ \mathbf{z}$. Then $g\mathbf{v}_i^2 \mathbf{y}\mathbf{C}\mathbf{z}\mathbf{D} = g\mathbf{y}\mathbf{v}_i^2 \mathbf{C}\mathbf{z}\mathbf{D}$.

In all the cases, the order of f is decreased while preserving $\mathbf{v}_i^2$. By induction on the order we get the conclusion. $\square$

Proposition 8 suggests a “square-free normalization” of vector-variable polynomials, by moving all squares to a set free of any reduction operation, and maintaining the set of squares in a normal form.

In a vector-variable monomial, let the set of squares be separated from the remainder of the monomial by a symbol “ $\square$ ”, such that all elements on the right side of the symbol are squares. Two things need to be established before such a symbol can be used in algebraic manipulations: (1) algebraic structure of the new symbolic system, (2) connection with the canonical system based on V2, V3, V4.

Let $\mathcal{S}$ be a commutative monoid. All elements in $\mathcal{S}$ span a $\mathbb{K}$ -vector space whose dimension equals the number of elements in $\mathcal{S}$. The product in the vector space is the multilinear extension of the product in $\mathcal{S}$. The vector space equipped with this product forms a commutative $\mathbb{K}$ -algebra, called the $\mathbb{K}$ -algebra extension of monoid $\mathcal{S}$, denoted by $\mathbb{K}\mathcal{S}$.

For a $\mathbb{K}$ -algebra $\mathcal{A}$, when $\mathcal{S}$ is a subset of the center of $\mathcal{A}$, then $\mathcal{A}$ is not only a module over the ring $\mathbb{K}\mathcal{S}$, but a multilinear algebra over $\mathbb{K}\mathcal{S}$, called a $\mathbb{K}\mathcal{S}$ -algebra.

For $\mathbb{K}$ -tensor algebra $\otimes[\mathcal{M}]$, let

$$\otimes^\square[\mathcal{M}] := \otimes[\mathcal{M}] / \langle \mathbf{V2} \rangle. \quad (5.1)$$

It is easy to see that when setting $\mathcal{S}$ to be generated by elements of the form $\mathbf{v}_i^2 := \mathbf{v}_i \otimes \mathbf{v}_i$, for all $\mathbf{v}_i \in \mathcal{M}$, then $\otimes^\square[\mathcal{M}]$ is a $\mathbb{K}\mathcal{S}$ -algebra, called the $\mathbb{K}\mathcal{S}$ -tensor algebra over multiset $\mathcal{M}$, or the *square-free tensor algebra* over $\mathcal{M}$. The product in $\otimes^\square[\mathcal{M}]$ is induced from the tensor product. For brevity we still denote the product by “ $\otimes$ ”, but denote the commutative product in $\mathcal{S}$ by juxtaposition of elements.

In $\otimes^\square[\mathcal{M}]$, for all $q \in \otimes[\mathcal{M}]$ and $s \in \mathcal{S}$, we introduce the notations

$$\begin{aligned} q\square s &:= q \otimes s \in \otimes[\mathcal{M}], \\ q\square &:= q \in \otimes[\mathcal{M}], \\ \square s &:= s \in \mathcal{S}. \end{aligned} \quad (5.2)$$

Then

$$(q_1\square s_1) \otimes (q_2\square s_2) := q_1 \otimes q_2\square s_1 s_2. \quad (5.3)$$

Formally, an element $q \in \mathcal{Q}[\mathcal{M}]$ is taken as $q\square 1$, and an element $s \in \mathbb{K}\mathcal{S}$ is taken as $1\square s$. In other words, factor $\square 1$ (or $\square$) in $q\square 1$ (or $q\square$) is usually omitted. So

$$q\square s = q \otimes (\square s) = (\square s) \otimes q, \quad (5.4)$$

and

$$\square st = (\square s) \otimes (\square t) = (\square t) \otimes (\square s). \quad (5.5)$$

That $\mathcal{S}$ is generated by squares can be succinctly expressed by the following identity:

$$\mathbf{v}_i \otimes \mathbf{v}_i = \square \mathbf{v}_i^2. \quad (5.6)$$

making left multiplication with f and right multiplication with g on both sides of the identity, we get $f \otimes (\mathbf{v}_i \otimes \mathbf{v}_i) \otimes g =$

$f \otimes g \square \mathbf{v}_i^2$. It includes V2 as a special case.

The *degree*, or *length*, of a monomial in $\otimes^\square[\mathcal{M}]$ is the degree of the monomial when taken as an element in $\otimes[\mathcal{M}]$. The *left degree* or *left length* of a monomial refers to the degree of the monomial on the left side of the square symbol. For a monomial $f \in \otimes^\square[\mathcal{M}]$, its *canonical form* in $\otimes[\mathcal{M}]$ is defined to be the monomial of lowest lexicographic order among all monomials equal to f modulo V2. The *order* of f is that of its canonical form. This ordering is still called the *lexicographic ordering*.

The canonical form of $f = \mathbf{v}_{i_1} \otimes \mathbf{v}_{i_2} \otimes \cdots \otimes \mathbf{v}_{i_k} \square \mathbf{v}_{j_1}^{2r_1} \mathbf{v}_{j_2}^{2r_2} \cdots \mathbf{v}_{j_l}^{2r_l}$, where $\mathbf{v}_{j_1} \prec \mathbf{v}_{j_2} \prec \cdots \prec \mathbf{v}_{j_l}$, can be obtained as follows:

1. Set $g = \mathbf{v}_{i_1} \otimes \mathbf{v}_{i_2} \otimes \cdots \otimes \mathbf{v}_{i_k}$.
2. For p from 1 to l , let $\mathbf{v}_{i_t}$ be the first variable in the sequence of g such that $\mathbf{v}_{i_t} \succ \mathbf{v}_{j_p} \succeq \mathbf{v}_{i_{t-1}}$. Insert

$$\underbrace{\mathbf{v}_{j_p} \otimes \mathbf{v}_{j_p} \otimes \cdots \otimes \mathbf{v}_{j_p}}_{2r_p}$$

to the position before $\mathbf{v}_{i_t}$ in g , and update g .

3. Output g .

The vector-variable polynomial ring $\mathcal{Q}[\mathcal{M}]$ when taken as the quotient of $\otimes^\square[\mathcal{M}]$ modulo the two-sided ideal $\mathcal{I}^\square[\mathcal{M}]$ generated by V3, V4, is a $\mathbb{K}S$ -algebra, called the *square-free polynomial ring*, denoted by $\mathcal{Q}^\square[\mathcal{M}]$. The product in $\mathcal{Q}^\square[\mathcal{M}]$ is still denoted by juxtaposition of elements. $\mathcal{I}^\square[\mathcal{M}]$ is called the *syzygy ideal* of $\mathcal{Q}^\square[\mathcal{M}]$.

Theorem 3 has the following square-free version for multiset $\mathcal{M}$:

THEOREM 9. *Let $\mathcal{M}$ be a multiset of m symbols, among which n are different ones: $\mathbf{v}_1 \prec \mathbf{v}_2 \prec \cdots \prec \mathbf{v}_n$, and let $\mathcal{I}^\square[\mathcal{M}]$ be the syzygy ideal of the square-free polynomial ring $\mathcal{Q}^\square[\mathcal{M}]$.*

(1) [Gröbner base] *The following are a reduced Gröbner base of $\mathcal{I}^\square[\mathcal{M}]$: G3, and*

Gj: *for all $3 < j < n+1$, and $i_1 < i_2 < \cdots < i_j$,*

$$[\mathbf{v}_{i_3} \mathbf{v}_{i_2} \mathbf{v}_{i_4} \mathbf{v}_{i_5} \cdots \mathbf{v}_{i_j} \mathbf{v}_{i_1}] - [\mathbf{v}_{i_2} \mathbf{v}_{i_4} \mathbf{v}_{i_5} \cdots \mathbf{v}_{i_j} \mathbf{v}_{i_1} \mathbf{v}_{i_3}],$$

EGk: *for all $3 \leq k \leq n+1$, and $i_1 < i_2 < \cdots < i_k$,*

$$[\mathbf{v}_{i_3} \mathbf{v}_{i_2} \mathbf{v}_{i_3} \mathbf{v}_{i_4} \cdots \mathbf{v}_{i_k} \mathbf{v}_{i_1}] - [\mathbf{v}_{i_2} \mathbf{v}_{i_3} \mathbf{v}_{i_4} \cdots \mathbf{v}_{i_k} \mathbf{v}_{i_1} \mathbf{v}_{i_3}].$$

The above Gröbner base is denoted by $\mathcal{G}^\square[\mathcal{M}]$.

(2) [Normal form] *In a normal form, every term is up to coefficient of the form $\mathbf{V}_{Y_1} \mathbf{v}_{z_1} \mathbf{V}_{Y_2} \mathbf{v}_{z_2} \cdots \mathbf{V}_{Y_k} \mathbf{v}_{z_k} \square s$ or*

$\mathbf{V}_{Y_1} \mathbf{v}_{z_1} \cdots \mathbf{V}_{Y_k} \mathbf{v}_{z_k} \mathbf{V}_{Y_{k+1}} \square s$, where

(i) $k \geq 0$,

(ii) $\mathbf{v}_{z_1} \mathbf{v}_{z_2} \cdots \mathbf{v}_{z_k}$ *is non-descending,*

(iii) *every $\mathbf{V}_{Y_i}$ is an ascending monomial of length > 0 ,*

(iv) $\mathbf{V}_{Y_1} \mathbf{V}_{Y_2} \cdots \mathbf{V}_{Y_k}$ (or $\mathbf{V}_{Y_1} \cdots \mathbf{V}_{Y_k} \mathbf{V}_{Y_{k+1}}$ if $\mathbf{V}_{Y_{k+1}}$ occurs) *is non-descending,*

(v) *for every $i \leq k$, let $\mathbf{v}_{t_i}$ be the trailing variable of $\mathbf{V}_{Y_i}$,*

then $\mathbf{v}_{t_i} \succ \mathbf{v}_{z_i}$,

(vi) *s is either 1 or the product of several squares.*

In $\mathcal{Q}^\square[\mathcal{M}]$, a polynomial is said to be $\mathcal{I}^\square$ -*normal* if its leading term is reduced with respect to the Gröbner base $\mathcal{G}^\square[\mathcal{M}]$.

A monic *square-free uni-bracket monomial* is of the form $[\mathbf{A}] \square s$, where $\mathbf{A}$ is either 1 or a monomial of length > 1 , s is a product of squares, and the length of $\mathbf{A}s$ is m . A *square-free uni-bracket polynomial* is a $\mathbb{K}$ -linear combination of square-free uni-bracket monomial. The space of square-free uni-bracket polynomials is denoted by $[\mathcal{Q}]^\square[\mathcal{M}]$.

The $\mathbb{K}$ -linear space of degree- m square-free polynomials is denoted by $\mathcal{Q}_m^\square[\mathcal{M}]$. The space $[\mathcal{Q}]^\square[\mathcal{M}]$ can be taken as the quotient of $\mathcal{Q}_m^\square[\mathcal{M}]$ modulo the ideal

$$\mathcal{J}^\square[\mathcal{M}] := \mathcal{I}^\square[\mathcal{M}] + [\mathcal{I}]^\square[\mathcal{M}], \quad (5.7)$$

where $[\mathcal{I}]^\square[\mathcal{M}]$ is composed of the vector parts of degree- m square-free polynomials, *i.e.*, the $\mathbb{K}$ -linear span of elements of the form

$$\mathbf{R}^\square : \mathbf{A} \square s - (-1)^a \mathbf{A}^\dagger \square s, \quad (5.8)$$

where $\mathbf{A}$ is a monomial of length $a > 0$ and contains no square, s is a product of squares, and the length of $\mathbf{A}s$ is m .

The modulo- $[\mathcal{I}]^\square[\mathcal{M}]$ operation identifies a square-free uni-bracket with its representative. It removes the outer bracket symbol on the left side of the “ $\square$ ” symbol from every square-free uni-bracket, disregarding the length of the bracket. Ideal $[\mathcal{I}]^\square[\mathcal{M}]$ is called the *uni-bracket removal ideal* in $\mathcal{Q}^\square[\mathcal{M}]$, and ideal is called the *syzygy ideal* of $[\mathcal{Q}]^\square[\mathcal{M}]$ in $\otimes^\square[\mathcal{M}]$.

6. GRÖBNER BASE AND NORMAL FORM FOR UNI-BRACKET POLYNOMIALS

In this section, we extend Theorem 7 to the case of general multiset $\mathcal{M}$ with $m \geq 3$ different vector variables. The modulo- $[\mathcal{I}]^\square[\mathcal{M}]$ operation is always assumed, *i.e.*, $[\mathcal{Q}]^\square[\mathcal{M}]$ and $\mathcal{Q}_m^\square[\mathcal{M}]$ are identical, and a uni-bracket does not have the outer bracket symbol on the left side of the “ $\square$ ” symbol.

The following are elements of $\mathcal{J}^\square[\mathcal{M}]$:

$$\begin{aligned} \mathbf{R}^\square(k) : & (\mathbf{K} - (-1)^k \mathbf{K}^\dagger) \square s, \\ \mathbf{S}_1(k) : & (\mathbf{i} \mathbf{A} \mathbf{b}_1 \mathbf{B} - \mathbf{b}_1 \mathbf{B} \mathbf{i} \mathbf{A}) \square s, \text{ for } \mathbf{i} \neq \mathbf{b}_1, \\ \mathbf{S}_{10}(k) : & (\mathbf{i} \mathbf{C} \mathbf{b}_1 - \mathbf{b}_1 \mathbf{i} \mathbf{C}) \square s, \text{ for } \mathbf{i} \neq \mathbf{b}_1, \\ \mathbf{S}_{11}(k) : & (\mathbf{b}_1 \mathbf{A} \mathbf{b}_1 \mathbf{B} - \mathbf{b}_1 \mathbf{B} \mathbf{b}_1 \mathbf{A}) \square s, \\ \mathbf{S}_{q1}(k) : & (\mathbf{b}_1 \mathbf{C} \mathbf{b}_1 - \mathbf{C} \square \mathbf{b}_1^2) \square s, \\ \mathbf{S}_{q1}[*](k) : & (\mathbf{b}_1 \mathbf{E} \mathbf{b}_1 - \mathbf{E} \square \mathbf{b}_1^2) [\mathbf{F}] \square s, \\ \mathbf{R}_1(k) : & \mathbf{b}_1 [\mathbf{D}] \square s, \\ \mathbf{R}_{11}(k) : & \mathbf{b}_1 [\mathbf{C} \mathbf{b}_1] \square s, \\ \mathbf{R}_{11}[*](k) : & \mathbf{b}_1 [\mathbf{E} \mathbf{b}_1] [\mathbf{F}] \square s, \\ \mathbf{R}_{12}(k) : & \mathbf{b}_1 [\mathbf{C} \mathbf{b}_2] \square s, \\ \mathbf{R}_{12}[*](k) : & \mathbf{b}_1 [\mathbf{E} \mathbf{b}_2] [\mathbf{F}] \square s, \end{aligned} \quad (6.1)$$

where

(a) s has length $m - k$, and the left length of each expression is $k > 0$;

(b) $\mathbf{b}_1, \mathbf{b}_2$ are respectively the variables of the lowest order and the second lowest order on the left side of the square symbol;

- (c) in S_1 , either $\mathbf{A}$ or $\mathbf{B}$ can be empty, while in S_{11} , both $\mathbf{A}$ and $\mathbf{B}$ are non-empty;
(d) in R_1 , $\mathbf{D}$ is either empty (i.e., $\mathbf{D} = 1$), or of length > 1 ;
(e) in S_{10} and R_{11} , $\mathbf{C}$ is non-empty;
(f) in $S_{q_1}[*](k)$ and $R_{11}[*]$, $\mathbf{E}, \mathbf{F}$ are non-empty, and $\mathbf{F}$ does not contain $\mathbf{b}_1$;
(g) in R_{12} , $\mathbf{C}$ is non-empty and does not contain $\mathbf{b}_1$;
(h) in $R_{12}[*]$, $\mathbf{E}, \mathbf{F}$ are non-empty and do not contain $\mathbf{b}_1$, and $\mathbf{F}$ does not contain $\mathbf{b}_2$.

In bracket $[\mathbf{1A1}]$, we have $[\mathbf{1A1}] \stackrel{\mathcal{I}^\square}{=} [\mathbf{A}]\square\mathbf{1}^2$. That the leading variable has higher order than the trailing variable is always possible. This is taken as a postulate for all the brackets in (6.1).

Consider a general element $f = (\mathbf{K} - (-1)^k \mathbf{K}^\dagger)\square s$ of type $R^\square(k)$:

1. If $\mathbf{b}_1$ occurs in $\mathbf{K}$ both as the leading variable and trailing variable, then $f \in \langle S_{q_1}(k), R^\square(k-2) \rangle$.
2. If $\mathbf{b}_1$ occurs in $\mathbf{K}$ at only one end, then $f \in \langle R_1(k), S_1(k) \rangle$.
3. If $\mathbf{b}_1$ occurs at the interior of $\mathbf{K}$, set $\mathbf{K} = \mathbf{Ab}_1\mathbf{B}$, where $\mathbf{A}, \mathbf{B}$ are both non-empty, and $\mathbf{b}_1$ does not occur at any end of $\mathbf{A}$ or $\mathbf{B}$. By (4.2), $f \in \langle R_1(k), S_1(k) \rangle$.

By induction on k , we get

LEMMA 10.

$$R^\square(k) \subseteq \sum_{h \leq k} (\langle S_1(h) \rangle + \langle S_{q_1}(h) \rangle + \langle R_1(h) \rangle).$$

In $R_1(k)$, when $\mathbf{D}$ contains $\mathbf{b}_1$, let $\mathbf{D} = \mathbf{Ab}_1\mathbf{B}$, where the lengths of $\mathbf{A}, \mathbf{B}$ are respectively $a, k - a - 2$, then

$$\begin{aligned} \mathbf{b}_1[\mathbf{Ab}_1\mathbf{B}]\square s &\stackrel{\mathcal{I}}{=} \mathbf{b}_1[\mathbf{BAAb}_1]\square s \\ &= (\mathbf{b}_1\mathbf{BAAb}_1 - (-1)^k \mathbf{A}^\dagger \mathbf{B}^\dagger)\square \mathbf{b}_1^2 s \\ &= (\mathbf{b}_1\mathbf{BAAb}_1 - \mathbf{BA}\square \mathbf{b}_1^2)\square s \\ &\quad + (\mathbf{BA} - (-1)^k \mathbf{A}^\dagger \mathbf{B}^\dagger)\square \mathbf{b}_1^2 s \\ &\in \langle S_{q_1}(k), R^\square(k-2) \rangle. \end{aligned} \quad (6.2)$$

By induction on k , we get that both $\mathbf{b}_1[\mathbf{Ab}_1\mathbf{B}]\square s$ and $\mathbf{b}_1[\mathbf{BAAb}_1]\square s \in R_{11}(k)$ are equivalent to $S_{q_1}(k)$: $(\mathbf{b}_1\mathbf{BAAb}_1 - \mathbf{BA}\square \mathbf{b}_1^2)\square s$ in the sense that their difference is in the ideal $\sum_{h \leq k-2} (\langle S_1(h) \rangle + \langle R_{11}(h) \rangle + \langle R_{12}(h) \rangle) + \langle R_1(1) \rangle + \mathcal{I}^\square[\mathcal{M}]$.

LEMMA 11. $R^\square(k)$ is a subset of the ideal

$$\sum_{h \leq k} (\langle S_1(h) \rangle + \langle R_{11}(h) \rangle + \langle R_{12}(h) \rangle) + \langle R_1(1) \rangle + \mathcal{I}^\square[\mathcal{M}].$$

THEOREM 12. Let $\mathcal{M}$ be a multiset of $m > 2$ symbols, among which $n \geq 2$ are different ones: $\mathbf{1} \prec \mathbf{2} \prec \dots \prec \mathbf{n}$, and let $\mathcal{Q}^\square[\mathcal{M}]$ be the square-free vector-variable polynomial ring over $\mathcal{M}$. Let $[\mathcal{Q}^\square[\mathcal{M}]]$ be the space of square-free uni-bracket polynomials, and let $\mathcal{J}^\square[\mathcal{M}]$ be its syzygy ideal in $\otimes^\square[\mathcal{M}]$.

(1) [Gröbner base] The following are a reduced Gröbner base of $\mathcal{J}^\square[\mathcal{M}]$, denoted by $\mathcal{BG}[\mathcal{M}]$:

$\mathcal{G}^\square[\mathcal{M}]$: Gi for all $3 \leq i \leq n$; EGj for all $3 \leq j \leq n+1$.

$S_1^\square$: $(\mathbf{Ab}_1\mathbf{B} - \mathbf{b}_1\mathbf{BA})\square s$, where the length of each term is m , $\mathbf{b}_1$ is the variable of the lowest order on the left side of the square symbol, $\mathbf{A}$ is a non-empty ascending sequence not containing $\mathbf{b}_1$, and $\mathbf{Ab}_1\mathbf{B}$ is $\mathcal{I}^\square$ -normal;

$R_1^\square[j, l]$:

$$\mathbf{b}_1[\mathbf{Y}_1\mathbf{b}_1][\mathbf{Y}_2\mathbf{b}_1] \cdots [\mathbf{Y}_j\mathbf{b}_1][\mathbf{Y}_{j+1}\mathbf{z}_{j+1}][\mathbf{Y}_{j+2}\mathbf{z}_{j+2}] \cdots [\mathbf{Y}_{j+l}\mathbf{z}_{j+l}]\square s, \quad (6.3)$$

where (i) $j, l \geq 0$, and the length of each term is m ,

(ii) $\mathbf{b}_1$ is the variable of the lowest order on the left side of the square symbol,

(iii) each $\mathbf{Y}_i$ is a non-empty ascending sequence not containing $\mathbf{b}_1$,

(iv) for all $j+1 \leq i \leq j+l$, $\mathbf{z}_i \neq \mathbf{b}_1$, and $\mathbf{z}_i\mathbf{Y}_i$ is ascending;

(v) $\mathbf{b}_1\mathbf{Y}_1\mathbf{b}_1 \cdots \mathbf{Y}_j\mathbf{b}_1\mathbf{Y}_{j+1}\mathbf{z}_{j+1} \cdots \mathbf{Y}_{j+l}\mathbf{z}_{j+l}$ is $\mathcal{I}^\square$ -normal.

(2) [Normal form] In a normal form, every term is $\mathcal{I}^\square$ -normal, and is up to coefficient of one of the following forms, where statements (ii), (iii) on $\mathbf{b}_1$ and the $\mathbf{Y}_i$ are still valid:

(I) $\mathbf{b}_1\mathbf{Y}_1\mathbf{b}_1 \cdots \mathbf{Y}_j\mathbf{b}_1\mathbf{Y}_{j+1}\mathbf{z}_{j+1} \cdots \mathbf{Y}_{j+l}\mathbf{z}_{j+l}\mathbf{Y}_{j+l+1}\square s$, where $j, l \geq 0$, and for all $j+1 \leq i \leq j+l$, $\mathbf{z}_i \prec \mathbf{t}_i$, the latter being the trailing variable of $\mathbf{Y}_i$;

(II) $\mathbf{b}_1\mathbf{Y}_1\mathbf{b}_1 \cdots \mathbf{Y}_j\mathbf{b}_1\mathbf{Y}_{j+1}\mathbf{z}_{j+1} \cdots \mathbf{Y}_{j+l}\mathbf{z}_{j+l}\square s$, where $j, k \geq 0$ but $j+k > 0$, each $\mathbf{z}_i \prec \mathbf{t}_i$, but for some $1 \leq h \leq l$, if $\mathbf{l}_{j+h}$ is the leading variable of $\mathbf{Y}_{j+h}$, then $\mathbf{l}_{j+h} \preceq \mathbf{z}_{j+h}$.

Remark. The set of $R_1^\square$ can be replaced by the following three sets of degree- m polynomials: for all $j > 0, l \geq 0$,

$$R_1^\square[0, 0] : \mathbf{b}_1\square s,$$

$$S_{q_1}^\square[j, l] : (\mathbf{b}_1\mathbf{Y}_1\mathbf{b}_1 - \mathbf{Y}_1\square \mathbf{b}_1^2)[\mathbf{Y}_2\mathbf{b}_1] \cdots [\mathbf{Y}_j\mathbf{b}_1][\mathbf{Y}_{j+1}\mathbf{z}_{j+1}] \cdots [\mathbf{Y}_{j+l}\mathbf{z}_{j+l}]\square s,$$

$$R_{12}^\square[j, l] : \mathbf{b}_1[\mathbf{Y}_1\mathbf{b}_2][\mathbf{Y}_2\mathbf{b}_2] \cdots [\mathbf{Y}_j\mathbf{b}_2][\mathbf{Y}_{j+1}\mathbf{z}_{j+1}][\mathbf{Y}_{j+2}\mathbf{z}_{j+2}] \cdots [\mathbf{Y}_{j+l}\mathbf{z}_{j+l}]\square s. \quad (6.4)$$

In $R_{12}^\square[j, l]$, $\mathbf{Y}_{j+1}, \dots, \mathbf{Y}_{j+l}$ do not contain $\mathbf{b}_2$, and the $\mathbf{z}_i \succ \mathbf{b}_2$. The replacement has no effect upon the normal forms.

Proof. There are several steps.

Step 1. We need to prove by induction on k that $S_1(k)$, $R_{11}(k)$, $R_{12}(k)$ are all in the ideal

$$\sum_{\text{left length} \leq k} \langle S_1^\square, R_1^\square[*] \rangle + \mathcal{I}^\square[\mathcal{M}], \quad (6.5)$$

where the asterisk stands for the (j, l) .

Once this is done, then since the leader of any element of type $S_1^\square$ or $R_1^\square[*]$ is $\mathcal{I}^\square$ -normal and cannot be cancelled by the leader of any other element of type $S_1^\square$ or $R_1^\square[*]$, the $S_1^\square, R_1^\square[*]$ and $\mathcal{G}^\square[\mathcal{M}]$ must be a reduced Gröbner base of $\langle S_1, R_{11}, R_{12}, R_1(1) \rangle + \mathcal{I}^\square[\mathcal{M}]$. By Lemma 11, this ideal is just $\mathcal{J}^\square[\mathcal{M}]$. This proves conclusion (1), and conclusion (2) follows.

Step 2. $S_1(k)$ requires $k > 1$; $R_{11}(k)$ and $R_{12}(k)$ both require $k > 2$. When $k = 2$, $S_1(2) = (\mathbf{b}_2\mathbf{b}_1 - \mathbf{b}_1\mathbf{b}_2)\square s$ is in $S_1^\square$. When $k = 3$, $R_{11}(3) = (\mathbf{b}_1\mathbf{b}_2\mathbf{b}_1 - \mathbf{b}_2\square\mathbf{b}_1^2)\square s = R_1^\square[1, 0]$, and $R_{12}(3) = \mathbf{b}_1[\mathbf{b}_3\mathbf{b}_2]\square s = R_1^\square[0, 1]$, where $\mathbf{b}_3 \succ \mathbf{b}_2$.

Consider $S_1(3)$. There are 3 elements led by variable $\mathbf{b}_2$: $(\mathbf{b}_2\mathbf{b}_1\mathbf{b}_2 - \mathbf{b}_1\square\mathbf{b}_2^2)\square s$, $(\mathbf{b}_2\mathbf{b}_1\mathbf{b}_3 - \mathbf{b}_1\mathbf{b}_3\mathbf{b}_2)\square s$, and $(\mathbf{b}_2\mathbf{b}_3\mathbf{b}_1 - \mathbf{b}_1\mathbf{b}_2\mathbf{b}_3)\square s$. They all belong to $S_1^\square$. There are two other elements in $S_1(3)$: $(\mathbf{b}_3\mathbf{b}_1\mathbf{b}_2 - \mathbf{b}_1\mathbf{b}_2\mathbf{b}_3)\square s$ and $(\mathbf{b}_3\mathbf{b}_2\mathbf{b}_1 - \mathbf{b}_1\mathbf{b}_3\mathbf{b}_2)\square s$. By

$$\begin{aligned} \mathbf{b}_3\mathbf{b}_2\mathbf{b}_1 - \mathbf{b}_1\mathbf{b}_3\mathbf{b}_2 &\stackrel{G_3}{=} \mathbf{b}_1\mathbf{b}_2\mathbf{b}_3 - \mathbf{b}_2\mathbf{b}_3\mathbf{b}_1; \\ \mathbf{b}_3\mathbf{b}_1\mathbf{b}_2 - \mathbf{b}_1\mathbf{b}_2\mathbf{b}_3 &\stackrel{G_3}{=} (\mathbf{b}_2\mathbf{b}_3\mathbf{b}_1 - \mathbf{b}_1\mathbf{b}_2\mathbf{b}_3) \\ &\quad + (\mathbf{b}_2\mathbf{b}_1\mathbf{b}_3 - \mathbf{b}_1\mathbf{b}_3\mathbf{b}_2), \end{aligned}$$

both are in $\langle S_1^\square \rangle + \mathcal{I}^\square[\mathcal{M}]$.

So the statement in Step 1 holds for $k \leq 3$. Assume that it holds for all $k < h$. When $k = h$, we need to make $\mathcal{I}^\square$ -reduction to the leaders of the elements of any of the types

$$S_1(h), R_{11}(h), R_{12}(h), R_{11}[*](h), R_{12}[*](h), \quad (6.6)$$

at the same time keep the reduction result to be within the $\mathbb{K}$ -linear space spanned by elements of the types listed in (6.6) but where the left length h is replaced by all $i \leq h$. Then ultimately all the leaders of these elements become $\mathcal{I}^\square$ -normal.

Step 3. Consider types $R_{11}(h), R_{11}[*](h), R_{12}(h), R_{12}[*](h)$. Let there be an $R_{11}(h)$ -typed element $f = 2(\mathbf{b}_1[\mathbf{A}\mathbf{b}_1]\square s)$, and an $R_{12}(h)$ -typed element $g = 2(\mathbf{b}_1[\mathbf{B}\mathbf{b}_2]\square s)$, where $\mathbf{B}$ does not contain $\mathbf{b}_1$. In the following we omit the factor $\square s$.

Do $\mathcal{I}^\square$ -reductions to $\mathbf{A}, \mathbf{B}$, and assume that the results are

$$\begin{aligned} \mathbf{A} &\stackrel{\mathcal{I}^\square}{=} \mathbf{C}_A\mathbf{b}_1 + \mathbf{b}_1\mathbf{D}_A + \mathbf{b}_1\mathbf{E}_A\mathbf{b}_1 + \mathbf{A}^N, \\ \mathbf{B} &\stackrel{\mathcal{I}^\square}{=} \mathbf{C}_B\mathbf{b}_2 + \mathbf{b}_2\mathbf{D}_B + \mathbf{b}_2\mathbf{E}_B\mathbf{b}_2 + \mathbf{B}^N, \end{aligned} \quad (6.7)$$

where (i) none of the terms in $\mathbf{C}_A, \mathbf{D}_A, \mathbf{E}_A, \mathbf{A}^N$ has $\mathbf{b}_1$ at any end;
(ii) none of the terms in $\mathbf{C}_B, \mathbf{D}_B, \mathbf{E}_B, \mathbf{B}^N$ has $\mathbf{b}_2$ at any end;
(iii) any of the four terms in each result may not occur;
(iv) the component on the right side of the square symbol in each term, together with the symbol itself, are omitted, as they do not affect the analysis below;
(v) in the extreme case, $\mathbf{A}^N$ or $\mathbf{B}^N$ may be in $\mathbb{K}$, if all vector variables in the term form squares and are moved to the right side of the square symbol.

Substituting the reduction results into f, g , we get

$$\begin{aligned} f &\stackrel{\mathcal{I}^\square}{=} (\mathbf{D}_A\mathbf{b}_1 - (-1)^h\mathbf{D}_A^\dagger\mathbf{b}_1)\square\mathbf{b}_1^2 && \in \langle S_1(h-2), R_1(h-2) \rangle \\ &\quad + (\mathbf{b}_1\mathbf{C}_A - (-1)^h\mathbf{b}_1\mathbf{C}_A^\dagger)\square\mathbf{b}_1^2 && \in R_1(h-2) \\ &\quad + (\mathbf{E}_A\square\mathbf{b}_1^2 - (-1)^h\mathbf{b}_1\mathbf{E}_A^\dagger\mathbf{b}_1)\square\mathbf{b}_1^2 && \in R_{11}(h-2) \\ &\quad + 2(\mathbf{b}_1[\mathbf{A}^N\mathbf{b}_1]), && \in R_{11}(h) \\ g &\stackrel{\mathcal{I}^\square}{=} 2(\mathbf{b}_1[\mathbf{C}_B])\square\mathbf{b}_2^2 && \in R_1(h-2) \\ &\quad + 2(\mathbf{b}_1[\mathbf{D}_B])\square\mathbf{b}_2^2 && \in R_1(h-2) \\ &\quad + 2(\mathbf{b}_1[\mathbf{E}_B\mathbf{b}_2])\square\mathbf{b}_2^2 && \in R_{12}(h-2) \\ &\quad + 2(\mathbf{b}_1[\mathbf{B}^N\mathbf{b}_2]). && \in R_{12}(h) \end{aligned}$$

Notice that the left lengths indicated on the right column are the maximal possible ones for the corresponding types. So

by induction hypothesis, we can assume that in f, g , monomials $\mathbf{A} = \mathbf{A}^N, \mathbf{B} = \mathbf{B}^N$ and both are $\mathcal{I}^\square$ -normal.

The $\mathcal{I}^\square$ -reduction to the leaders of f, g are much the same with the procedure in the proof of Theorem 7 starting from Step 3 there to Step 8, with negligible revisions. Formula (4.11) can also be used to split the leader of factor $\mathbf{b}_1[\mathbf{Y}_1\mathbf{b}_1 \cdots \mathbf{Y}_k\mathbf{b}_1]$ in a type- $R_{11}[*]$ element, and the leader of factor $\mathbf{b}_1[\mathbf{Y}_1\mathbf{b}_2 \cdots \mathbf{Y}_k\mathbf{b}_2]$ in a type- $R_{12}[*]$ element.

By induction on the order of the leader, we get that $R_{11}(h), R_{12}(h), \text{Sq}_1(h), \text{Sq}_1[*](h), R_{11}[*](h), R_{12}[*](h)$ are all in (6.5) where $k = h$.

Step 4. Consider a general type- $S_{10}(h)$ element $f = (\mathbf{A}\mathbf{b}_1 - \mathbf{b}_1\mathbf{A})\square s$. Let the $\mathcal{I}^\square$ -reduction result of $\mathbf{A}$ be as in (6.7). Then if omitting “ $\square s$ ”,

$$\begin{aligned} f &\stackrel{\mathcal{I}^\square}{=} \begin{array}{ll} \mathbf{C}_A\square\mathbf{b}_1^2 - \mathbf{b}_1\mathbf{C}_A\mathbf{b}_1 & \in \text{Sq}_1(h) \\ + \mathbf{b}_1\mathbf{D}_A\mathbf{b}_1 - \mathbf{D}_A\square\mathbf{b}_1^2 & \in \text{Sq}_1(h) \\ + (\mathbf{b}_1\mathbf{E}_A - \mathbf{E}_A\mathbf{b}_1)\square\mathbf{b}_1^2 & \in S_1(h-2) \\ + \mathbf{A}^N\mathbf{b}_1 - \mathbf{b}_1\mathbf{A}^N & \in S_{10}(h) \end{array} \end{aligned}$$

So we can assume that in $f = (\mathbf{A}\mathbf{b}_1 - \mathbf{b}_1\mathbf{A})\square s$, monomial $\mathbf{A} = \mathbf{A}^N$ and is $\mathcal{I}^\square$ -normal.

The $\mathcal{I}^\square$ -reduction to the leading term of f is much the same with the procedure in the proof of Lemma 5 starting from Case (i) there to Case (ii). By induction on the order of the leading term, we get that $S_{10}(h)$ is in (6.5) where $k = h$.

Step 5. Consider a general type- $S_1(h)$ element $g = (\mathbf{A}\mathbf{b}_1\mathbf{B} - \mathbf{b}_1\mathbf{B}\mathbf{A})\square s$, where $\mathbf{A}, \mathbf{B}$ are both non-empty. Let the $\mathcal{I}^\square$ -reduction results of $\mathbf{A}, \mathbf{B}$ be as in (6.7), where every $\mathbf{b}_2$ is replaced by $\mathbf{b}_1$. Then if omitting “ $\square s$ ”,

$$\begin{aligned} g &\stackrel{\mathcal{I}^\square}{=} \begin{array}{l} \mathbf{C}_A\mathbf{C}_B\mathbf{b}_1\square\mathbf{b}_1^2 - \mathbf{b}_1\mathbf{C}_B\mathbf{b}_1\mathbf{C}_A\mathbf{b}_1 \\ + (\mathbf{C}_A\mathbf{b}_1\mathbf{D}_B - \mathbf{D}_B\mathbf{C}_A\mathbf{b}_1)\square\mathbf{b}_1^2 \\ + (\mathbf{C}_A\mathbf{b}_1\mathbf{E}_B\mathbf{b}_1 - \mathbf{E}_B\mathbf{b}_1\mathbf{C}_A\mathbf{b}_1)\square\mathbf{b}_1^2 \\ + \mathbf{C}_A\mathbf{B}^N\square\mathbf{b}_1^2 - \mathbf{b}_1\mathbf{B}^N\mathbf{C}_A\mathbf{b}_1 \\ + \mathbf{b}_1\mathbf{D}_A\mathbf{b}_1\mathbf{C}_B\mathbf{b}_1 - \mathbf{b}_1\mathbf{C}_B\mathbf{D}_A\square\mathbf{b}_1^2 \\ + (\mathbf{b}_1\mathbf{D}_A\mathbf{D}_B - \mathbf{D}_B\mathbf{b}_1\mathbf{D}_A)\square\mathbf{b}_1^2 \\ + \mathbf{b}_1\mathbf{D}_A\mathbf{E}_B\mathbf{b}_1\square\mathbf{b}_1^2 - \mathbf{E}_B\mathbf{D}_A\square\mathbf{b}_1^4 \\ + \mathbf{b}_1\mathbf{D}_A\mathbf{b}_1\mathbf{B}^N - \mathbf{b}_1\mathbf{B}^N\mathbf{b}_1\mathbf{D}_A \\ + (\mathbf{b}_1\mathbf{E}_A\mathbf{C}_B\mathbf{b}_1 - \mathbf{b}_1\mathbf{C}_B\mathbf{E}_A\mathbf{b}_1)\square\mathbf{b}_1^2 \\ + (\mathbf{b}_1\mathbf{E}_A\mathbf{b}_1\mathbf{D}_B - \mathbf{D}_B\mathbf{b}_1\mathbf{E}_A\mathbf{b}_1)\square\mathbf{b}_1^2 \\ + \mathbf{b}_1\mathbf{E}_A\mathbf{b}_1\mathbf{E}_B\mathbf{b}_1\square\mathbf{b}_1^2 - \mathbf{E}_B\mathbf{E}_A\mathbf{b}_1\square\mathbf{b}_1^4 \\ + \mathbf{b}_1\mathbf{E}_A\mathbf{B}^N\mathbf{b}_1^2 - \mathbf{b}_1\mathbf{B}^N\mathbf{b}_1\mathbf{E}_A\mathbf{b}_1 \\ + \mathbf{A}^N\mathbf{b}_1\mathbf{C}_B\mathbf{b}_1 - \mathbf{b}_1\mathbf{C}_B\mathbf{b}_1\mathbf{A}^N \\ + (\mathbf{A}^N\mathbf{D}_B - \mathbf{D}_B\mathbf{A}^N)\square\mathbf{b}_1^2 \\ + \mathbf{A}^N\mathbf{b}_1\mathbf{E}_B\mathbf{b}_1 - \mathbf{b}_1\mathbf{E}_B\mathbf{b}_1\mathbf{A}^N \\ + \mathbf{A}^N\mathbf{b}_1\mathbf{B}^N - \mathbf{b}_1\mathbf{B}^N\mathbf{A}^N. \end{array} \end{aligned}$$

In the above result, the lines that do not belong to the ideal $\langle \text{Sq}_1(h), S_1(h-2), \text{Sq}_1(h-2), S_1(h-4) \rangle$ are

$$\begin{aligned} &\mathbf{b}_1\mathbf{D}_A\mathbf{b}_1\mathbf{B}^N - \mathbf{b}_1\mathbf{B}^N\mathbf{b}_1\mathbf{D}_A && \in \langle S_{11}(h), \text{Sq}_1(h) \rangle \\ &+ \mathbf{A}^N\mathbf{b}_1\mathbf{C}_B\mathbf{b}_1 - \mathbf{b}_1\mathbf{C}_B\mathbf{b}_1\mathbf{A}^N && \in \langle S_{11}(h), S_{10}(h) \rangle \\ &+ \mathbf{A}^N\mathbf{b}_1\mathbf{E}_B\mathbf{b}_1 - \mathbf{b}_1\mathbf{E}_B\mathbf{b}_1\mathbf{A}^N && \in \langle S_{11}(h), S_{10}(h) \rangle \\ &+ \mathbf{A}^N\mathbf{b}_1\mathbf{B}^N - \mathbf{b}_1\mathbf{B}^N\mathbf{A}^N && \in S_1(h) \end{aligned} \quad (6.8)$$

Some remarks on (6.8) are necessary. The first line of (6.8), if

nonzero, is $S_{11}(h)$ when $\mathbf{B}^N \notin \mathbb{K}$, and $S_{q_1}(h)$ otherwise. By $\mathbf{A}^N \mathbf{b}_1 \mathbf{C}_B \mathbf{b}_1 - \mathbf{b}_1 \mathbf{C}_B \mathbf{b}_1 \mathbf{A}^N = (\mathbf{A}^N \mathbf{b}_1 \mathbf{C}_B \mathbf{b}_1 - \mathbf{b}_1 \mathbf{A}^N \mathbf{b}_1 \mathbf{C}_B) + (\mathbf{b}_1 \mathbf{A}^N \mathbf{b}_1 \mathbf{C}_B - \mathbf{b}_1 \mathbf{C}_B \mathbf{b}_1 \mathbf{A}^N)$, the second line of (6.8) is a $\mathbb{K}$ -linear combination of an element of type $S_{10}(h)$ and another element of type $S_{11}(h)$.

Consider a general type- $S_{11}(h)$ element $p = (\mathbf{b}_1 \mathbf{A} \mathbf{b}_1 \mathbf{B} - \mathbf{b}_1 \mathbf{B} \mathbf{b}_1 \mathbf{A}) \square s$, where the length of $\mathbf{A}$ is $a > 0$. When omitting $\square s$,

$$\begin{aligned} p &\stackrel{\mathcal{I}^\square}{=} \mathbf{b}_1 \mathbf{B} (\mathbf{b}_1 \mathbf{A} - (-1)^a \mathbf{A}^\dagger \mathbf{b}_1) \\ &\quad + (-1)^a \mathbf{A}^\dagger \mathbf{B} \square \mathbf{b}_1^2 - \mathbf{b}_1 \mathbf{B} \mathbf{b}_1 \mathbf{A} \\ &= (-1)^a (\mathbf{A}^\dagger \mathbf{B} \square \mathbf{b}_1^2 - \mathbf{b}_1 \mathbf{B} \mathbf{A}^\dagger \mathbf{b}_1) \\ &\stackrel{S_{q_1}(h)}{=} (-1)^a (\mathbf{A}^\dagger \mathbf{B} - \mathbf{B} \mathbf{A}^\dagger) \square \mathbf{b}_1^2 \\ &\in \langle S_1(h-2) \rangle. \end{aligned}$$

So $S_{11}(h)$ is in (6.5) where $k = h$.

By (6.8), we can assume that in $g = (\mathbf{A} \mathbf{b}_1 \mathbf{B} - \mathbf{b}_1 \mathbf{B} \mathbf{A}) \square s$, monomials $\mathbf{A} = \mathbf{A}^N$, $\mathbf{B} = \mathbf{B}^N$ and both are $\mathcal{I}^\square$ -normal. The $\mathcal{I}^\square$ -reduction to the leading term of g is much the same with that in the proof of Lemma 5 starting from Case (i) there to Case (iii). By induction on the order of the leading term, we get that $S_1(h)$ is in (6.5) where $k = h$. $\square$

Consider a bracket polynomial f whose multiset of variables is $\mathcal{M}$. In any term of f , when all the brackets but one are expanded into two terms by definition, f is changed into a uni-bracket polynomial g . Using the Gröbner base $\mathcal{BG}[\mathcal{M}]$ to make reduction to g results in a uni-bracket polynomial h , where each term is $\mathcal{I}^\square$ -normal. h must have the lowest order lexicographically among all uni-bracket polynomials equal to f . It is called the *lowest-representative normal form* of g , or the *uni-bracket normal form* of f .

Remark. In the above definition of normal forms, we only considered square-free ones. Of course any normal form can be converted to a canonical one, where all representatives are $\mathcal{I}$ -normal instead of $\mathcal{I}^\square$ -normal. Later on, we consider only square-free ones.

Let the size of $\mathcal{M}$ be m . Given any partition $(i_1, \dots, i_k)$ of integer m , where each $i_j > 1$, there is a *Caianiello expansion* [6] of uni-bracket polynomials into bracket polynomials where each term is composed of k brackets of length $i_1, \dots, i_k$ respectively. Each expansion produces a normal form. It is not clear if such a normal form is of any value.

7. NORMALIZATION OF BRACKET POLYNOMIALS

In this section, we do NOT remove bracket symbols from uni-brackets in multiset of variables $\mathcal{M}$.

Consider attaching an additional vector variable $\mathbf{v}_0 \notin \mathcal{M}$ to $\mathcal{M}$ to form a bigger multiset $\tilde{\mathcal{M}}$. Let $\mathbf{v}_0 \prec$ all variables in $\mathcal{M}$. In the procedure of obtaining $R_1^\square$ by the $\mathcal{I}^\square$ -reduction of $\mathbf{b}_1[\mathbf{A}]$ in the proof of Theorem 12, or in more details, in the proof of Theorem 7, if the input is $\mathbf{v}_0[\mathbf{A}]$ where $\mathbf{A} \in \mathcal{Q}_m[\mathcal{M}]$, then among the Gröbner base of $\mathcal{I}^\square[\tilde{\mathcal{M}}]$, only those elements in $\mathcal{G}^\square[\mathcal{M}]$ are needed in $\mathcal{I}^\square$ -reduction.

The reduction of $\mathbf{v}_0[\mathbf{A}]$ is a procedure of recursively doing $\mathcal{I}^\square$ -reductions to the leader of the bracket polynomial obtained from previous $\mathcal{I}^\square$ -reductions to $[\mathbf{A}]$. At any instance, the representative of a bracket monomial in reduction is the leader of the bracket monomial. The reduction results in a $\mathbb{K}$ -linear combination of monomials of the form $\mathbf{v}_0[\mathbf{Y}_1 \mathbf{z}_1] \cdots [\mathbf{Y}_k \mathbf{z}_k]$, where $\mathbf{z}_i \mathbf{Y}_i$ is ascending for every i . After removing $\mathbf{v}_0$ from the result, we get another normal form of uni-bracket $[\mathbf{A}]$ in $\mathcal{M}$.

THEOREM 13. *Let f be a square-free bracket polynomial in multiset of variables $\mathcal{M}$. Do the following to f :*

1. *Always select the leader of a bracket as its representative.*
2. *Rearrange the order of the brackets in the same term, so that the leading variables of the brackets are non-descending.*
3. *Use (4.6) to normalize the interior of a bracket; it also splits a bracket into two.*
4. *Use (4.9) to absorb a bracket into the one ahead of it.*
5. *Use (4.10) to decrease the order of the product of two brackets by lifting a lower-order variable from the second bracket to the first.*
6. *Use (4.11) to segment a long bracket of type $\mathbf{Y}_1 \mathbf{z}_1 \mathbf{Y}_2 \mathbf{z}_2 \cdots \mathbf{Y}_k \mathbf{z}_k$ into short ones.*
7. *Once the representative of the leading term of f is $\mathcal{I}^\square$ -normal, output the leading term, and continue the above $\mathcal{I}^\square$ -normalization to the remainder of f .*

The output, called the lowest-leader normal form, or leader-normal form is a bracket polynomial where the representative of each term is up to coefficient of the form $[\mathbf{Y}_1 \mathbf{z}_1][\mathbf{Y}_2 \mathbf{z}_2] \cdots [\mathbf{Y}_k \mathbf{z}_k] \square s$, where each $\mathbf{z}_i \mathbf{Y}_i$ is ascending, $\mathbf{Y}_1 \mathbf{Y}_2 \cdots \mathbf{Y}_k$ and $\mathbf{z}_1 \mathbf{z}_2 \cdots \mathbf{z}_k$ are both non-descending, and the leading variable $\mathbf{l}_i$ of each $\mathbf{Y}_i$ satisfies $\mathbf{l}_i \succ \mathbf{z}_i$. Two bracket polynomials are equal if and only if their leader-normal forms are identical.

Proof. After operations 1 and 2, for any bracket monomial in the reduction procedure, its representative is also its leader, so that the representative of the leading term of bracket polynomial f is the leader of f . Once the leader of f is $\mathcal{I}^\square$ -normal, it is the leading term of the normal form of vector-variable polynomial f with respect to the Gröbner base $\mathcal{G}^\square[\mathcal{M}]$. By induction on the order of the output terms from the highest down, we get the uniqueness of the leader-normal form for f .

For two equal bracket polynomials, they have identical uni-bracket normal forms, and so have identical leader-normal forms. $\square$

From the above proof, we see that the leader-normal form of a bracket polynomial f has the following properties: (1) the representative of any term is the leader of the term; (2) the representative of the leading term is the leading term of the $\mathcal{I}^\square$ -normal form of vector-variable polynomial f .

By Theorem 12, the Gröbner base $\mathcal{BG}[\tilde{\mathcal{M}}]$ of $\mathcal{I}^\square[\tilde{\mathcal{M}}]$ is composed of $\mathcal{G}^\square[\mathcal{M}]$ and the $R_1^\square[*] : \mathbf{v}_0 g \square s$ for all monomials g in the leader-normal forms of bracket polynomials in $\mathcal{M}$, such that g has length m . This phenomenon is easy to understand: the $\mathbb{K}$ -linear subspace of $\mathcal{I}^\square[\tilde{\mathcal{M}}]$ composed of polynomials whose terms are led by variable $\mathbf{v}_0$, is the space

of degree- $(m+1)$ polynomials of the form $\mathbf{v}_0 f \square s$, for all bracket polynomials f in $\mathcal{M}$ such that fs has length m . The leader-normal forms are a basis of the $\mathbb{K}$ -linear space of length- m square-free bracket polynomials in $\mathcal{M}$.

In a leader-normal form, if we commute $\mathbf{Y}_i$ and $\mathbf{z}_i$ in $[\mathbf{Y}_i \mathbf{z}_i]$, we get another normal form whose terms are up to coefficient of the form $[\mathbf{z}_1 \mathbf{Y}_1] \cdots [\mathbf{z}_k \mathbf{Y}_k] \square s$. If we write the left side of the square symbol in the following tableau form, where $\mathbf{Y}_i = \mathbf{y}_{i1} \mathbf{y}_{i2} \cdots \mathbf{y}_{it_i}$, we get

$$\begin{bmatrix} \mathbf{z}_1 & \mathbf{y}_{11} & \cdots & \cdots & \mathbf{y}_{1t_1} \\ \mathbf{z}_2 & \mathbf{y}_{21} & \cdots & \mathbf{y}_{2t_2} & \\ \vdots & \vdots & \ddots & & \\ \mathbf{z}_k & \mathbf{y}_{k1} & \cdots & \cdots & \mathbf{y}_{kt_k} \end{bmatrix},$$

- where (1) each row does not need to have equal length, and it is not required that the length be non-increasing as in Young tableau;
(2) each row is an ascending sequence of variables;
(3) each column is a non-descending sequence of variables;
(4) $\mathbf{y}_{it_i} \preceq \mathbf{y}_{(i+1)1}$ for $1 \leq i < k$.

Such a normal form is called the *straight form*. Feature (4) above makes this definition stronger than the *straight form* (or *standard form*) of Young tableau. In comparison, in classical bracket algebra a bracket monomial is in straight form if and only if the entries are ascending along each row, and non-descending along each column.

The procedure of deriving the straight form of a bracket polynomial is called *straightening*. Among the formulas used in Theorem 13 for straightening, (4.10) is highly nontrivial and requires further investigation.

Set $\mathbf{B}\mathbf{u}$, $\mathbf{a}\mathbf{D}$ in (4.10) to be new $\mathbf{A}, \mathbf{B}$ respectively, and let the lengths of $\mathbf{B}, \mathbf{C}$ be b, c . Then (4.10) can be written succinctly as follows:

$$\begin{aligned} [\mathbf{A}\mathbf{v}][\mathbf{B}\mathbf{w}\mathbf{C}] &= [\mathbf{A}\mathbf{w}\mathbf{C}\mathbf{v}][\mathbf{B}] - (-1)^b [\mathbf{A}\mathbf{w}\mathbf{B}^\dagger \mathbf{v}][\mathbf{C}] \\ &\quad - (-1)^b [\mathbf{w}\mathbf{C}\mathbf{v}][\mathbf{A}\mathbf{B}^\dagger] + (-1)^b [\mathbf{w}\mathbf{B}^\dagger \mathbf{v}][\mathbf{A}\mathbf{C}] \\ &\quad - [\mathbf{A}\mathbf{w}][\mathbf{C}\mathbf{v}\mathbf{B}]. \end{aligned} \quad (7.1)$$

It is called the *shuffle formula for bracket normalization*.

PROPOSITION 14. For any two monomials $\mathbf{A}, \mathbf{B}$ of length a, b respectively,

$$\begin{cases} \frac{\mathbf{AB} + \mathbf{BA}}{2} = [\mathbf{AB}] + (-1)^b (\mathbf{A}[\mathbf{B}^\dagger] - [\mathbf{A}]\mathbf{B}^\dagger), \\ \frac{\mathbf{AB} - (-1)^{a+b} \mathbf{A}^\dagger \mathbf{B}^\dagger}{2} = (-1)^a ([\mathbf{A}^\dagger]\mathbf{B} - \mathbf{A}^\dagger[\mathbf{B}]). \end{cases} \quad (7.2)$$

Proof.

$$\begin{aligned} \mathbf{AB} + \mathbf{BA} &= 2[\mathbf{AB}] - (-1)^{a+b} \mathbf{B}^\dagger \mathbf{A}^\dagger + 2[\mathbf{B}]\mathbf{A} - (-1)^b \mathbf{B}^\dagger \mathbf{A} \\ &= 2([\mathbf{AB}] + [\mathbf{B}]\mathbf{A} - (-1)^b \mathbf{B}^\dagger [\mathbf{A}]). \end{aligned}$$

□

The fundamental $\mathcal{I}$ -reduction formula (4.5) is a direct consequence of the first identity in (7.2) for $\mathbf{AB} = \mathbf{u}\mathbf{D}$. The

shuffle formula (7.1) is a consequence of the following identity by making left multiplication with $\mathbf{A}$ and then applying the bracket operator to both sides of the identity:

$$\begin{aligned} \mathbf{v}[\mathbf{B}\mathbf{w}\mathbf{C}] + \mathbf{w}[\mathbf{C}\mathbf{v}\mathbf{B}] &= (-1)^b (\mathbf{w}\mathbf{C}\mathbf{v}[\mathbf{B}^\dagger] - [\mathbf{w}\mathbf{C}\mathbf{v}]\mathbf{B}^\dagger \\ &\quad - \mathbf{w}\mathbf{B}^\dagger \mathbf{v}[\mathbf{C}] + [\mathbf{w}\mathbf{B}^\dagger \mathbf{v}]\mathbf{C}). \end{aligned} \quad (7.3)$$

The identity can be obtained as follows: by the second identity of (7.2) from right to left, the right side of (7.3) equals

$$2^{-1} (\mathbf{B}\mathbf{w}\mathbf{C}\mathbf{v} + \mathbf{v}\mathbf{B}\mathbf{w}\mathbf{C}) - (-1)^{b+c} 2^{-1} (\mathbf{B}^\dagger \mathbf{v}\mathbf{C}^\dagger \mathbf{w} + \mathbf{w}\mathbf{B}^\dagger \mathbf{v}\mathbf{C}^\dagger),$$

which by the first equality of (7.2), equals

$$\begin{aligned} &[\mathbf{v}\mathbf{B}\mathbf{w}\mathbf{C}] - (-1)^{b+c} \mathbf{v}[\mathbf{C}^\dagger \mathbf{w}\mathbf{B}^\dagger] \\ &\quad - (-1)^{b+c} [\mathbf{w}\mathbf{B}^\dagger \mathbf{v}\mathbf{C}^\dagger] + \mathbf{w}[\mathbf{C}\mathbf{v}\mathbf{B}] \\ &= \mathbf{v}[\mathbf{B}\mathbf{w}\mathbf{C}] + \mathbf{w}[\mathbf{C}\mathbf{v}\mathbf{B}]. \end{aligned}$$

The shuffle formula can be further generalized. In monomial $[\mathbf{A}\mathbf{v}\mathbf{D}][\mathbf{B}\mathbf{w}\mathbf{C}]$, let the leading variable and trailing variable of any sequence $\mathbf{F}$ be $\mathbf{l}_F$ and $\mathbf{t}_F$ respectively. Assume $\mathbf{l}_A \preceq \mathbf{l}_B$, and $\mathbf{l}_A \succ \mathbf{t}_D$, and $\mathbf{l}_B \succ \mathbf{t}_C$. Further assume $\mathbf{l}_A \succ \mathbf{v} \succ \mathbf{w}$. Then $\mathcal{I}^\square$ -reductions can be made to $[\mathbf{A}\mathbf{v}\mathbf{D}][\mathbf{B}\mathbf{w}\mathbf{C}]$ to decrease its leader, leading to the following result:

PROPOSITION 15. Let the lengths of monomials $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$ be a, b, c, d respectively. Then

$$\begin{aligned} [\mathbf{A}\mathbf{v}\mathbf{D}][\mathbf{B}\mathbf{w}\mathbf{C}] &= [\mathbf{v}\mathbf{D}\mathbf{B}\mathbf{w}][\mathbf{A}\mathbf{C}] - (-1)^{b+c} [\mathbf{v}\mathbf{D}\mathbf{C}^\dagger \mathbf{w}][\mathbf{A}\mathbf{B}^\dagger] \\ &\quad - (-1)^d [\mathbf{A}\mathbf{w}][\mathbf{D}^\dagger \mathbf{v}\mathbf{B}\mathbf{C}] - [\mathbf{v}\mathbf{D}][\mathbf{A}\mathbf{w}\mathbf{B}\mathbf{C}] \\ &\quad - (-1)^b [\mathbf{A}\mathbf{w}\mathbf{B}^\dagger \mathbf{v}\mathbf{D}][\mathbf{C}] + [\mathbf{A}\mathbf{w}\mathbf{C}\mathbf{v}\mathbf{D}][\mathbf{B}]. \end{aligned} \quad (7.4)$$

Proof.

$$\begin{aligned} &4[\mathbf{A}\mathbf{v}\mathbf{D}][\mathbf{B}\mathbf{w}\mathbf{C}] \\ &= \mathbf{A}\mathbf{v}\mathbf{D}\mathbf{B}\mathbf{w}\mathbf{C} - (-1)^{b+c} \mathbf{A}\mathbf{v}\mathbf{D}\mathbf{C}^\dagger \mathbf{w}\mathbf{B}^\dagger \\ &\quad - (-1)^{a+d} \mathbf{D}^\dagger \mathbf{v}\mathbf{A}^\dagger \mathbf{B}\mathbf{w}\mathbf{C} + (-1)^{a+b+c+d} \mathbf{D}^\dagger \mathbf{v}\mathbf{A}^\dagger \mathbf{C}^\dagger \mathbf{w}\mathbf{B}^\dagger \\ &\stackrel{\mathcal{I}^\square}{=} (\mathbf{v}\mathbf{D}\mathbf{B}\mathbf{w} + (-1)^{b+d} \mathbf{w}\mathbf{B}^\dagger \mathbf{D}^\dagger \mathbf{v}) \mathbf{A}\mathbf{C} \\ &\quad - (-1)^b ((-1)^c \mathbf{v}\mathbf{D}\mathbf{C}^\dagger \mathbf{w} + (-1)^d \mathbf{w}\mathbf{D}\mathbf{C}^\dagger \mathbf{v}) \mathbf{A}\mathbf{B}^\dagger \\ &\quad - (-1)^{b+d} \mathbf{A}\mathbf{w}\mathbf{B}^\dagger \mathbf{D}^\dagger \mathbf{v}\mathbf{C} + (-1)^{b+d} \mathbf{A}\mathbf{w}\mathbf{C}\mathbf{D}^\dagger \mathbf{v}\mathbf{B}^\dagger \\ &\quad - (-1)^{a+d} \mathbf{D}^\dagger \mathbf{v}(\mathbf{B}\mathbf{w}\mathbf{C} - (-1)^{b+c} \mathbf{C}^\dagger \mathbf{w}\mathbf{B}^\dagger) \mathbf{A}^\dagger \\ &\stackrel{\mathcal{I}^\square}{=} 4([\mathbf{v}\mathbf{D}\mathbf{B}\mathbf{w}][\mathbf{A}\mathbf{C}] - (-1)^{b+c} [\mathbf{v}\mathbf{D}\mathbf{C}^\dagger \mathbf{w}][\mathbf{A}\mathbf{B}^\dagger]) \\ &\quad + (-1)^{a+d} (\mathbf{B}\mathbf{C}\mathbf{D}^\dagger \mathbf{v} - \mathbf{D}^\dagger \mathbf{v}\mathbf{B}\mathbf{C}) \mathbf{w}\mathbf{A}^\dagger \\ &\quad + (-1)^{b+d} \mathbf{A}\mathbf{w}(\mathbf{C}\mathbf{D}^\dagger \mathbf{v}\mathbf{B}^\dagger - \mathbf{B}^\dagger \mathbf{D}^\dagger \mathbf{v}\mathbf{C}). \end{aligned} \quad (7.5)$$

By (7.2),

$$\begin{aligned} &(-1)^d (\mathbf{B}\mathbf{C}\mathbf{D}^\dagger \mathbf{v} - \mathbf{D}^\dagger \mathbf{v}\mathbf{B}\mathbf{C}) \\ &= (-1)^d \mathbf{B}\mathbf{C}\mathbf{D}^\dagger \mathbf{v} - (-1)^{b+c} \mathbf{v}\mathbf{D}\mathbf{C}^\dagger \mathbf{B}^\dagger \\ &\quad + (-1)^{b+c} (\mathbf{v}\mathbf{D} - (-1)^d \mathbf{D}^\dagger \mathbf{v}) \mathbf{C}^\dagger \mathbf{B}^\dagger \\ &\quad + (-1)^{c+d} \{ \mathbf{D}^\dagger \mathbf{v}(-[\mathbf{C}^\dagger]\mathbf{B} + \mathbf{C}^\dagger[\mathbf{B}]) \}, \end{aligned}$$

we get from (7.5) the following:

$$\begin{aligned} &4[\mathbf{A}\mathbf{v}\mathbf{D}][\mathbf{B}\mathbf{w}\mathbf{C}] \\ &\stackrel{\mathcal{I}^\square}{=} 4([\mathbf{v}\mathbf{D}\mathbf{B}\mathbf{w}][\mathbf{A}\mathbf{C}] - (-1)^{b+c} [\mathbf{v}\mathbf{D}\mathbf{C}^\dagger \mathbf{w}][\mathbf{A}\mathbf{B}^\dagger]) \\ &\quad + 4[\{(-1)^d [\mathbf{B}\mathbf{C}\mathbf{D}^\dagger \mathbf{v}] + (-1)^{b+c} [\mathbf{v}\mathbf{D}]\mathbf{C}^\dagger \mathbf{B}^\dagger \\ &\quad - (-1)^{c+d} \mathbf{D}^\dagger \mathbf{v}\mathbf{B}[\mathbf{C}^\dagger] + (-1)^{c+d} \mathbf{D}^\dagger \mathbf{v}\mathbf{C}^\dagger[\mathbf{B}]\} (-1)^a \mathbf{w}\mathbf{A}^\dagger \\ &\quad + (-1)^b \mathbf{A}\mathbf{w}\{(-1)^c \mathbf{v}\mathbf{D}\mathbf{C}^\dagger \mathbf{B}^\dagger + (-1)^c \mathbf{B}^\dagger \mathbf{C}^\dagger \mathbf{v}\mathbf{D} \\ &\quad + (-1)^d \mathbf{C}\mathbf{D}^\dagger \mathbf{v}\mathbf{B}^\dagger - (-1)^d \mathbf{B}^\dagger \mathbf{D}^\dagger \mathbf{v}\mathbf{C}\} \end{aligned}$$

$$\begin{aligned} \stackrel{\square}{=} & 4([\mathbf{vDBw}][\mathbf{AC}] - (-1)^{b+c}[\mathbf{vDC}^\dagger \mathbf{w}][\mathbf{AB}^\dagger]) \\ & -4\{(-1)^d[\mathbf{Aw}][\mathbf{BCD}^\dagger \mathbf{v}] + [\mathbf{AwBC}][\mathbf{vD}] \\ & + (-1)^b[\mathbf{AwB}^\dagger \mathbf{vD}][\mathbf{C}] - [\mathbf{AwCvD}][\mathbf{B}]\}. \end{aligned}$$

□

Clearly each term in the result of (7.4) has lower leader than the input. To better understand this reduction formula, we write it in tableau form:

$$\begin{aligned} \left[\begin{array}{cc} \mathbf{AvD} \\ \mathbf{BwC} \end{array} \right] &= \left[\begin{array}{cc} \mathbf{AwCvD} \\ \mathbf{B} \end{array} \right] + \left[\begin{array}{cc} \mathbf{A}((-1)^{b+1}\mathbf{Bw})^\dagger \mathbf{vD} \\ \mathbf{C} \end{array} \right] \\ &+ \left[\begin{array}{cc} \mathbf{vD} \\ ((-1)^b \mathbf{B}^\dagger) \mathbf{w} ((-1)^a \mathbf{A})^\dagger ((-1)^c \mathbf{C}^\dagger) \end{array} \right] \\ &+ \left[\begin{array}{cc} \mathbf{wC}((-1)^{d+1} \mathbf{vD})^\dagger \\ \mathbf{B}((-1)^a \mathbf{A})^\dagger \end{array} \right] \\ &+ \left[\begin{array}{cc} ((-1)^{b+1} \mathbf{Bw})^\dagger ((-1)^{d+1} \mathbf{vD})^\dagger \\ \mathbf{AC} \end{array} \right] \\ &+ \left[\begin{array}{cc} \mathbf{Aw} \\ ((-1)^b \mathbf{B}^\dagger) \mathbf{vD} ((-1)^c \mathbf{C}^\dagger) \end{array} \right]. \end{aligned} \quad (7.6)$$

As $\mathbf{l_A} \succ \mathbf{v} \succ \mathbf{w}$ and $\mathbf{l_A} \succ \mathbf{t_D}$, to decrease the order of the leader, in the first line of (7.6), a subsequence or reversed subsequence of the second row of the input tableau is moved up between $\mathbf{A}, \mathbf{v}$ of the first row, with the requirement that the subsequence be led by $\mathbf{w}$. In the second line of (7.6), $\mathbf{A}$ is moved down behind $\mathbf{w}$ of the second row.

In the third and fourth lines of (7.6), the leading subsequence $\mathbf{A}$ of the first row is commuted with a subsequence or reversed subsequence of the second row led by $\mathbf{w}$. In the last line of (7.6), the trailing subsequence $\mathbf{vD}$ of the first row is commuted with variable $\mathbf{w}$ of the second row.

We make comparison with the shuffle formula for straightening in classical bracket algebra [13]. In the classical bracket algebra over 3D vector space, where the exterior product is also denoted by juxtaposition of elements, suppose that

$$\left[\begin{array}{ccc} \mathbf{a} & \mathbf{v} & \mathbf{d} \\ \mathbf{b} & \mathbf{w} & \mathbf{c} \end{array} \right]$$

is not straight: $\mathbf{avd}$ is ascending, so is $\mathbf{bwc}$; $\mathbf{ab}$ is non-descending, but $\mathbf{v} \succ \mathbf{w}$. The shuffle formula is obtained as follows. For any four vectors $\mathbf{v}, \mathbf{d}, \mathbf{b}, \mathbf{w}$ of the 3D vector space, their exterior product equals zero. By

$$\begin{aligned} 0 &= \mathbf{a} \vee (\mathbf{vdbw}) \vee \mathbf{c} \\ &= [\mathbf{avd}][\mathbf{bwc}] - [\mathbf{avb}][\mathbf{dwc}] + [\mathbf{avw}][\mathbf{dbc}] \\ &\quad + [\mathbf{adb}][\mathbf{vwc}] - [\mathbf{adw}][\mathbf{vbc}] + [\mathbf{abw}][\mathbf{vdc}], \end{aligned} \quad (7.7)$$

where “ $\vee$ ” is the dual of the exterior product called the *meet product* [14], we get the following shuffle formula, also called *van der Waerden relation*:

$$\left[\begin{array}{ccc} \mathbf{a} & \mathbf{v} & \mathbf{d} \\ \mathbf{b} & \mathbf{w} & \mathbf{c} \end{array} \right] = \left[\begin{array}{ccc} \mathbf{a} & \mathbf{v} & \mathbf{b} \\ \mathbf{d} & \mathbf{w} & \mathbf{c} \end{array} \right] + \left[\begin{array}{ccc} \mathbf{a} & \mathbf{v} & \mathbf{w} \\ \mathbf{b} & \mathbf{d} & \mathbf{c} \end{array} \right]$$

$$\begin{aligned} &+ \left[\begin{array}{ccc} \mathbf{a} & \mathbf{b} & \mathbf{d} \\ \mathbf{v} & \mathbf{w} & \mathbf{c} \end{array} \right] + \left[\begin{array}{ccc} \mathbf{a} & \mathbf{w} & \mathbf{d} \\ \mathbf{b} & \mathbf{v} & \mathbf{c} \end{array} \right] \\ &- \left[\begin{array}{ccc} \mathbf{a} & \mathbf{b} & \mathbf{w} \\ \mathbf{v} & \mathbf{d} & \mathbf{c} \end{array} \right]. \end{aligned} \quad (7.8)$$

- (1) The first line commutes $\mathbf{d}$ of the first row and one of the first two vectors of the second row;
- (2) the second line commutes $\mathbf{v}$ of the first row and one of the first two vectors of the second row;
- (3) the last line commutes $\mathbf{vd}$ of the first row and the first two vectors of the second row.

For straightening in classical bracket algebra, there are other operations besides (7.8). When $\mathbf{a} \succ \mathbf{b}$, we only need to commute the two brackets. When $\mathbf{d} \succ \mathbf{c}$ while $\mathbf{a} \preceq \mathbf{b}$ and $\mathbf{v} \preceq \mathbf{w}$, we have another formula for straightening. By

$$\begin{aligned} 0 &= (\mathbf{av}) \vee (\mathbf{dbwc}) \\ &= [\mathbf{avd}][\mathbf{bwc}] - [\mathbf{avb}][\mathbf{dwc}] \\ &\quad + [\mathbf{avw}][\mathbf{dbc}] - [\mathbf{avc}][\mathbf{dbw}], \end{aligned}$$

we get the following *Grassmann-Plücker relation*:

$$\left[\begin{array}{ccc} \mathbf{a} & \mathbf{v} & \mathbf{d} \\ \mathbf{b} & \mathbf{w} & \mathbf{c} \end{array} \right] = \left[\begin{array}{ccc} \mathbf{a} & \mathbf{v} & \mathbf{b} \\ \mathbf{d} & \mathbf{w} & \mathbf{c} \end{array} \right] + \left[\begin{array}{ccc} \mathbf{a} & \mathbf{v} & \mathbf{w} \\ \mathbf{b} & \mathbf{d} & \mathbf{c} \end{array} \right] + \left[\begin{array}{ccc} \mathbf{a} & \mathbf{v} & \mathbf{c} \\ \mathbf{b} & \mathbf{w} & \mathbf{d} \end{array} \right].$$

The last vector $\mathbf{d}$ of the first row commutes in turn with every vector of the second row.

8. CONCLUSION

In the bottom-up approach to manipulating brackets, long brackets are expanded into basic ones by Caianiello expansion, in the end only brackets of length 2 and 3 are left for further algebraic manipulations. This approach proves to be inefficient in practice, despite the fact that there are straightening algorithms for polynomials of basic invariants [3], [10].

Uni-bracket polynomials provide a top-down approach to manipulating brackets. Given a bracket polynomial, by “ungrading”, each bracket but one in every term is expanded into a vector-variable binomial, and the bracket polynomial is changed into a uni-bracket one. Algebraic manipulations of uni-bracket polynomials can take full advantage of the associativity of the vector-variable product and the symmetries within a uni-bracket. The Gröbner base $\mathcal{G}^\square[\mathcal{M}]$ provided by this paper further fulfills the arsenal of symbolic manipulations on uni-bracket polynomials.

The last section of this paper suggests a third approach to manipulating brackets by algebraic manipulations directly upon the input brackets. To establish this approach there are many research topics ahead: division among bracket polynomials, properties of principal ideals, bracket polynomial factorization, and simplification by reducing the number of terms, *etc.* This seems to be a promising approach.

This paper is supported partially by NSFC 10871195, 60821002/F02, and NCMIS of CAS. The Gröbner bases for $m = 5, 6$ were first computed by Dr. L. Huang on his implementation of the classical non-commutative Gröbner base algorithm [9].

9. REFERENCES

- [1] Altmann, S.L. *Rotations, Quaternions, and Double Groups*. Oxford University Press, Oxford, 1986.
- [2] Bravi, P. and Brini, A. Remarks on invariant geometric calculus, Cayley-Grassmann algebras and geometric Clifford algebras. In: Crapo, H. and Senato, D. (eds.), *Algebraic Combinatorics and Computer Science*, Springer, Milano, pp. 129-150, 2001.
- [3] DeConcini, C., Eisenbud, E. and Procesi, C. Hodge algebras, *Astérisque* **91**, 1982.
- [4] Hestenes, D. and Sobczyk, G. *Clifford Algebra to Geometric Calculus*. D. Reidel, Dordrecht, Boston, 1984.
- [5] Li, H. A recipe for symbolic geometric computing: long geometric product, BREEFS and Clifford factorization. In: Brown, C.W. (ed.), *Proc. ISSAC 2007*, ACM Press, New York, pp. 261-268, 2007.
- [6] Li, H. *Invariant Algebras and Geometric Reasoning*. World Scientific, Singapore, 2008.
- [7] Li, H., Huang, L., and Liu, Y. Normalization of quaternionic polynomials. arxiv: 1301.5338v1 [math.RA], 2013.
- [8] Lounesto, P. *Clifford Algebras and Spinors*. Cambridge University Press, Cambridge, 1997.
- [9] Mora, T. An introduction to commutative and noncommutative Gröbner bases. *Theoretical Computer Science* **134** (1994) 131-173.
- [10] Mourrain, B. and Stolfi, N. Computational symbolic geometry. In: White, N. (ed.), *Invariant Methods in Discrete and Computational Geometry*, D. Reidel, Dordrecht, Boston, pp. 107-139, 1995.
- [11] Olver, P.J. *Classical Invariant Theory*. Cambridge University Press, Cambridge, 1999.
- [12] Sturmfels, B. Computing final polynomials and final syzygies using Buchberger's Gröbner basis method. *Result in Math.* **15**: 551-560, 1989.
- [13] Sturmfels, B. and White, N. Gröbner bases and invariant theory. *Advances in Math.* **76**: 245-259, 1989.
- [14] White, N. The bracket ring of combinatorial geometry I. *Trans. Amer. Math. Soc.* **202**: 79-103, 1975.